

**PATIENT**

Brody Hasbury

**SPECIES**

Canine

**BREED**

Norfolk Terrier

**SEX**

Neutered Male

**AGE**

9 Years

**WEIGHT**

6.7 kg

**INTERPRETED BY**Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)**IMAGING  
PERFORMED BY**

Dr. Gira

**HOSPITAL NAME**

Sabadilla Animal Clinic

**REFERRING VET**Dr. Nathaniel  
Asemadahun**INVOICE**

72063

**DATE**

1/8/26

**PRESENTING CLINICAL SIGNS**

Progressive ALT elevation, lower USG. Patient is not PUPD and otherwise asymptomatic  
Abnormal PE/Chem/CBC/UA Results: Attached

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN****Urinary System**

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The prostate is normal in size (0.51 cm) and shape for this neutered male dog. The parenchyma is homogenous and the external margins are smooth. The prostatic urethra appears normal with no evidence of irregularity, invasion, mass effect or calculi.

The left kidney has a normal shape and size (4.4 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (3.53 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

**Adrenal Glands**

The left adrenal gland is normal in size measuring 0.35 cm at the cranial pole and 0.45 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is normal in size measuring 0.45 cm at the cranial pole and 0.35 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

**Spleen**

The spleen is subjectively normal in size (1.3 cm), echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

**Liver**

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is homogenous echotexture. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

The gall bladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and likely incidental at this time. The cystic and common bile ducts are normal/not visible.



## PATIENT

Brody Hasbury

## SPECIES

Canine

## BREED

Norfolk Terrier

## SEX

Neutered Male

## AGE

9 Years

## WEIGHT

6.7 kg

## INTERPRETED BY

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

## IMAGING PERFORMED BY

Dr. Gira

## HOSPITAL NAME

Sabadilla Animal Clinic

## REFERRING VET

Dr. Nathaniel  
Asemadahun

## INVOICE

72063

## DATE

1/8/26

## *Gastrointestinal*

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. Duodenum wall measures 0.45 cm. Jejunum wall measures 0.33 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

## *Pancreas*

The pancreas is normal and isoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

## *Free Abdomen*

Evaluation of the peritoneal cavity did not reveal any evidence of effusion. There is no significant lymphadenopathy. A jejunal lymph node is visualized measuring 0.40 cm. The omentum is normal in echogenicity.

## ULTRASONOGRAPHIC FINDINGS

- No significant ultrasonographic lesions visualized.

## INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

No significant lesions are visualized associated with the liver. Unfortunately, there are many causes for an ALT elevation that cannot be diagnosed by ultrasound alone. Consider the following for further evaluation:

- Recommend pre- and post-prandial bile acids to assess liver function.
- Consider a repeat evaluation of bilirubin to confirm that it is elevated.
- If clinically appropriate, consider screening for Leptospirosis.

Further options could include a fine needle aspirate. This could evaluate for neoplastic causes, infectious, etc. It is less helpful for inflammatory or other hepatopathies, which would likely require biopsies for diagnosis. Prior to considering biopsy, you could consider treatment for acute liver injury with a course of Ursodiol, Denamarin, +/- antibiotics.

If ALT is persistently elevated, particularly if liver function is abnormal and/or the bilirubin continues to be elevated, biopsies of the liver should be considered with samples for histopathology, culture and copper levels.



**PATIENT**

Brody Hasbury

**SPECIES**

Canine

**BREED**

Norfolk Terrier

**SEX**

Neutered Male

**AGE**

9 Years

**WEIGHT**

6.7 kg

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

**IMAGING  
PERFORMED BY**

Dr. Gira

**HOSPITAL NAME**

Sabadilla Animal Clinic

**REFERRING VET**


Dr. Nathaniel  
Asemadahun

**INVOICE**

72063

**DATE**

1/8/26





## PATIENT

Brody Hasbury

## SPECIES

Canine

## BREED

Norfolk Terrier

## SEX

Neutered Male

## AGE

9 Years

## WEIGHT

6.7 kg

## INTERPRETED BY

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

## IMAGING PERFORMED BY

Dr. Gira

## HOSPITAL NAME

Sabadilla Animal Clinic

## REFERRING VET


Dr. Nathaniel  
Asemadahun

## INVOICE

72063

## DATE

1/8/26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

info@sonopath.com