

**PATIENT**

Lux Moran

**PRESENTING CLINICAL SIGNS**

**SPECIES**

Canine

No PU/PD Came in for annual exam/vaccines- LABS doneRS: 09-15-21 at 1:10p: CBC WNL Chem - moderate elevation ALP 612, new for her (last labs 2019 ALP 79) UA - USG 1.012 (isosthenuric), sed inactive, no protein T4 Normal 1.5 ALP - r/o benign hepatopathy, HAC, neoplasia Isosthenuria - r/o early renal dz vs incidental/hydration  
Abnormal PE/Chem/CBC/UA Results: mild sedation with Dexdormitor

**BREED**

Lab Retriever

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**SEX**

Spayed Female

**Urinary System**  
The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of mucosal irregularities, masses or cystic calculi. Bladder wall measured 0.37 cm. The proximal urethra appears somewhat prominent with a wall thickness measuring 0.39 cm, smooth with no obvious irregularities. This could just be anatomic variation, but correlate these findings with urinalysis findings (looking for hematuria) and for any evidence of straining. Correlate with rectal exam findings.

**AGE**

10 Years

The left kidney has a normal shape and size (6.76 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

**WEIGHT**

68 Pounds

The right kidney has a normal shape and size (6.4 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

**Adrenal Glands**

The left adrenal gland is normal in size measuring 0.71 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

**IMAGING PERFORMED BY**

Loetitia Saint-Jacques, RVT

The right adrenal gland is normal in size measuring 0.61 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

**HOSPITAL NAME**

Alpine AH

**Spleen**

The spleen is subjectively normal in size. The spleen echotexture is heterogenous and mottled, the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. There is a very small hypoechoic nodule measuring 0.46 cm x 0.37 cm in the parenchyma.

**REFERRING VET**

Dr. Rachel

**Liver**

**INVOICE**

26041

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is heterogenous in echotexture with subtle, indistinct focal mottling. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

**DATE**

10/1/21



**PATIENT**

Lux Moran The gallbladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are primarily anechoic. The cystic and common bile ducts are normal/not visible.

**SPECIES**

**Gastrointestinal**

Canine

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

**BREED**

Lab Retriever

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. The duodenum measured as normal (between 0.3-0.5cm in wall thickness) and the jejunum measured as normal (between 0.2-0.47cm.)

**SEX**

Spayed Female

Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

**AGE**

10 Years

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

**Pancreas**

**WEIGHT**

68 Pounds

The pancreas is normal and isoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

**Free Abdomen**

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

**Other**

**IMAGING PERFORMED BY**

Loetitia Saint-Jacques, RVT

No significant pericardial effusion.

**PRIMARY FINDINGS**

**HOSPITAL NAME**

Alpine AH

- Mildly heterogeneous liver – The diffuse hepatic changes are non-specific and could be consistent with vacuolar hepatopathy, nodular hyperplasia, inflammatory/immune-mediated disease, fibrosis, extramedullary hematopoiesis, toxic hepatopathy (e.g., copper), infiltrative neoplasia (less likely) or other hepatopathy.

**REFERRING VET**

Dr. Rachel

- Mildly mottled spleen with small, hypoechoic nodule – There is a non-cavitated, hypoechoic splenic nodule visualized. Differentials include lymphoid hyperplasia, extramedullary hematopoiesis, infiltrative neoplasia, inflammation, other. Cytology or histopathology would be necessary to get a definitive diagnosis.

**INVOICE**


26041

**SECONDARY FINDINGS**

- Subjectively thickened urethral wall – this is a subjective finding and may not be significant. Correlate with rectal exam findings, symptoms, and urinalysis results. If concerned, consider reevaluation.

**DATE**

10/1/21



**PATIENT**

Lux Moran

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

No focal hepatic lesions were observed, and the biliary tract appears relatively normal. These are factors I consider when evaluating a patient for an ALP elevation.

**SPECIES**

Canine

An elevation in ALP is a common finding. In general, however, causes of ALP elevation fall into three primary categories:

**BREED**

Lab Retriever

Induction phenomena, biliary diseases, and primary liver disorders.

**SEX**

Spayed Female

- Induction phenomena is the most common for elevation in ALP. These are systemic illnesses that 'turn on' the liver enzyme. Causes of this include Cushing's disease, dental disease, arthritis, and numerous others. In many cases the exact cause is unclear but as long as ultrasound and bile acids tests are normal most patients do not have progressive changes in their liver. While liver biopsy is not routinely performed, vacuolar hepatopathy, is noted on most biopsies. This is often non-progressive but in rare cases can be more severe and lead to liver failure.

**AGE**

10 Years

- If signs of cushings disease are present recommend endocrine function testing to evaluate for cushings disease.
- Consider fine needle aspirate to rule out round cell neoplasia -if this is a concern.

**WEIGHT**

68 Pounds

- If a cause for the ALP elevation is not identified: I recommend recheck general blood work every 6 months, ultrasound once per year, and bile acids test every 1-2 years based on other results. If the ALP continues to climb a biopsy could be considered.

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

- Consider long term use of denamarin, and monitoring for the signs of cushings developing.
- A primary vacuolar hepatopathy can be breed related and is seen in Scottish Terriers, Schnauzers, Cocker spaniels etc..

**IMAGING PERFORMED BY**

Loetitia Saint-Jacques, RVT

A small, hypoechoic splenic nodule is visualized. Options moving forward are fine needle aspirate of that area/nodule, or continued monitoring with ultrasound (recheck in 3-4 weeks).

**HOSPITAL NAME**

Alpine AH

The thickening described in the proximal urethra I suspect is normal variation, but correlate to any clinical signs or urinalysis changes that may make this more clinically relevant.

**REFERRING VET**


Dr. Rachel

**INVOICE**

26041

**DATE**

10/1/21





**PATIENT**

Lux Moran

**SPECIES**

Canine

**BREED**

Lab Retriever

**SEX**

Spayed Female

**AGE**

10 Years

**WEIGHT**

68 Pounds

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

**IMAGING PERFORMED BY**

Loetitia Saint-Jacques, RVT

**HOSPITAL NAME**

Alpine AH

**REFERRING VET**





Dr. Rachel


**INVOICE**

26041

**DATE**

10/1/21





**PATIENT**

Lux Moran

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

**SPECIES**

Canine

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**BREED**

Lab Retriever

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

kathleen.sennello@sonopath.com

**SEX**

Spayed Female

**AGE**

10 Years

**WEIGHT**

68 Pounds

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

**IMAGING PERFORMED BY**

Loetitia Saint-Jacques, RVT

**HOSPITAL NAME**

Alpine AH

**REFERRING VET**

Dr. Rachel

**INVOICE**

26041

**DATE**

10/1/21