

**PATIENT**

Hannah Durling

**SPECIES**

Feline

**BREED**

DSH

**SEX**

Spayed Female

**AGE**

14 Years

**WEIGHT**

3.8 kg

**INTERPRETED BY**

Kathleen Sennello  
DVM, MS, Diplomate  
ACVIM (Small Animal  
Internal Medicine)

**IMAGING PERFORMED BY**

Kelly Reschny

**HOSPITAL NAME**

Buck Animal Hospital

**REFERRING VET**

Dr. Sommers

**INVOICE**

26718

**DATE**

10/28/21

**PRESENTING CLINICAL SIGNS**

frequent vomiting, history of constipation, tiny hard stool palpable on exam, Grade 1-2 murmur currently on buprenorphine and ondansetron  
Abnormal PE/Chem/CBC/UA Results: august 2021 ALKP <10, Sept 2021 fPL abnormal

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**Urinary System**

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The left kidney is small in size at 1.76 cm and is rounded in shape. There is very little normal architecture visualized, as the kidney appears shrunken and appears to contain two nephroliths measuring 0.37 cm and 0.41 cm. There is no evidence of perinephric inflammation or effusion, and there is no evidence of pyelectasia, infarcts, or hydroureter.

The right kidney has a normal shape and size (4.14 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

**Adrenal Glands**

The left adrenal gland is normal in size measuring 0.34 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is normal in size measuring 0.39 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

**Spleen**

The spleen is subjectively normal in size, echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

**Liver**

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is homogenous echotexture. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

The gallbladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are primarily anechoic. The cystic and common bile ducts are normal/not visible.

**Gastrointestinal**

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.36cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.



**PATIENT**

Hannah Durling

The visualized areas of duodenum, jejunum and ileum have a uniform diameter with minimal fluid distension. Wall thickness is normal to slightly increased. Bowel loops follow a typical curvilinear path with distinct wall layering, but some areas display a prominent muscularis layer which does not display the typical 1:3 muscularis:mucosa layer ratio. Jejunum wall measured 0.23 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

**SPECIES**

Feline

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

**BREED**

DSH

**Pancreas**

The pancreas is normal and isoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

**SEX**

Spayed Female

**Free Abdomen**

Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

**AGE**

14 Years

**ULTRASONOGRAPHIC FINDINGS**

**WEIGHT**

3.8 kg

- Small left kidney with minimal architecture and nephroliths – This is likely a minimally functional kidney. This could have been a progressive change or a congenital issue.
- Prominent muscularis layer to the small intestine – The small intestinal wall changes are most consistent with an inflammatory process (i.e., inflammatory bowel disease) with a low possibility of emerging lymphoma. This can be a common finding in older cats.

**INTERPRETED BY**

Kathleen Sennello  
DVM, MS, Diplomate  
ACVIM (Small Animal  
Internal Medicine)

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

There are no focal lesions observed to explain the vomiting and constipation reported in the history. If not already performed, recommend abdominal radiographs to look for evidence of obstipation or any foreign material that was not evident on today's scan. If obstipated, the vomiting could be due to lack of progressive motility. If not obstipated, then vomiting could be due to a stomach or small intestinal issue.

**IMAGING PERFORMED BY**

Kelly Reschny

- Recommend current bloodwork including thyroid panel and GI panel with an fPLI, TLI, cobalamin and folate to further evaluate for pancreatic and small intestinal disease.
- Recommend starting a stool softener such as Miralax and ensuring the patient is adequately hydrated.
- If able, consider a diet trial with a hydrolyzed protein or novel protein diet.
- If symptoms are persisting or getting worse and there is no response to symptomatic therapy, then recommend endoscopy of the large and small bowel.

**REFERRING VET**


Dr. Sommers


**INVOICE**

26718

**DATE**

10/28/21





**PATIENT**

Hannah Durling

**SPECIES**

Feline

**BREED**

DSH

**SEX**

Spayed Female

**AGE**

14 Years

**WEIGHT**

3.8 kg

**INTERPRETED BY**

Kathleen Sennello  
DVM, MS, Diplomate  
ACVIM (Small Animal  
Internal Medicine)

**IMAGING  
PERFORMED BY**

Kelly Reschny

**HOSPITAL NAME**

Buck Animal Hospital

**REFERRING VET**


Dr. Sommers

**INVOICE**

26718

**DATE**

10/28/21



The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

kathleen.sennello@sonopath.com