



PATIENT

Zoe Goldsworthy

SPECIES

Feline

BREED

DSH

SEX

Spayed Female

AGE

6 Years 1 Month

WEIGHT

8 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

M. Kermendy, CVT

HOSPITAL NAME

Wauwatosa Veterinary
Clinic

REFERRING VET

Dr. Kevin Kicker

INVOICE

71743

DATE

11/12/25

PRESENTING CLINICAL SIGNS

4-week history of vomiting decreased appetite and now diarrhea in the last 3 days. Blood results have been unremarkable and CPI testing is normal. Owner has been unsuccessful in administering empiric treatments including cerenia and mirtaz. Presenting for abdominal ultrasound in an effort to isolate disease and level of severity for concise treatment planning. Unremarkable exam, approximately 19% decrease in body weight in 1 month.

Abnormal PE/Chem/CBC/UA Results: T4 normal on 10/20/25 QPL normal on 11/8/25 Very mild decrease in total protein but normal albumin and globulins

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The left kidney has a normal shape and size (3.35 cm) with a small cortical cyst measuring 0.43 cm and occasional pinpoint cortical mineralizations most consistent with dystrophic mineralization. Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (3.35 cm) with occasional pinpoint cortical mineralizations most consistent with dystrophic mineralization. Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, infarcts or hydroureter. Renal vasculature is normal.

Adrenal Glands

The left adrenal gland is normal in size measuring 0.38 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The region of the right adrenal (between right cranial kidney and vena cava) is unremarkable, but the adrenal is not distinctly visualized. No evidence of a mass effect is visualized.

Spleen

The spleen is subjectively normal in size (0.63 cm), echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

Liver

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is homogenous echotexture. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

**PATIENT**

Zoe Goldsworthy

The gall bladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and likely incidental at this time. The cystic and common bile ducts are normal/not visible.

SPECIES

Feline

Gastrointestinal

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.36cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

BREED

DSH

The visualized areas of duodenum, jejunum and ileum have a uniform diameter with minimal fluid distension. Wall thickness is normal to slightly increased. Bowel loops follow a typical curvilinear path with distinct wall layering, but some areas display a prominent muscularis layer which does not display the typical 1:3 muscularis:mucosa layer ratio. Duodenum wall measures 0.26 cm. Jejunum wall measures 0.24 cm. Visualized peristalsis appears appropriate. The muscularis layer appears diffusely prominent.

SEX

Spayed Female

AGE

6 Years 1 Month

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

WEIGHT

8 lbs

Pancreas

The pancreas is normal and isoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

Free Abdomen

Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

IMAGING PERFORMED BY

M. Kermendy, CVT

ULTRASONOGRAPHIC FINDINGS

- Mild inflammatory type pattern visualized associated with the small intestine (prominent muscularis).

HOSPITAL NAME

Wauwatosa Veterinary
Clinic

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The small intestine is somewhat prominent with a prominent muscularis layer. These changes are most consistent with mild chronic inflammatory type change (IBD type pattern). An early neoplastic change cannot be ruled out. Consider the following:

- Consider a novel protein/hydrolyzed protein diet (exclusively at least 4-6 weeks)
- Consider a GI panel to Texas A&M for evaluation of B12 levels, folate, PLI/TLI etc.. to further evaluate for pancreatic/small intestinal disease.
- Recommend chronic probiotic therapy.

INVOICE

71743

DATE

11/12/25

If symptoms are persistent despite taking these measures, consider obtaining biopsies of the GI tract for further evaluation. Additionally, repeat imaging in the future could be considered, looking for the progression of today's lesion or the development of new lesions.



PATIENT

Zoe Goldsworthy

SPECIES

Feline

BREED

DSH

SEX

Spayed Female

AGE

6 Years 1 Month

WEIGHT

8 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

M. Kermendy, CVT

HOSPITAL NAME

Wauwatosa Veterinary
Clinic

REFERRING VET


Dr. Kevin Kicker

INVOICE

71743

DATE

11/12/25





PATIENT

Zoe Goldsworthy

SPECIES

Feline

BREED

DSH

SEX

Spayed Female

AGE

6 Years 1 Month

WEIGHT

8 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

M. Kermendy, CVT

HOSPITAL NAME

Wauwatosa Veterinary
Clinic

REFERRING VET


Dr. Kevin Kicker

INVOICE

71743

DATE

11/12/25



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

info@sonopath.com