



PATIENT

Mama Agricola

SPECIES

Feline

BREED

DSH

SEX

Spayed Female

AGE

9 years

WEIGHT

12.5 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Pamela Harrigan, RDCS,
Certified Veterinary
Sonographer

HOSPITAL NAME

Norfolk County
Veterinary Service

REFERRING VET

Dr. Jeremy Carignan

INVOICE

10751

DATE

11/13/2025

PRESENTING CLINICAL SIGNS

Chronic diarrhea, weight loss, history of mildly elevated kidney values in the past. CBC/Chem/UA pending.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The left kidney has a normal shape and size (4.03 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (3.84 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

Adrenal Glands

The left adrenal gland is normal in size measuring 0.33 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is normal in size measuring 0.36 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

Spleen

The spleen is subjectively normal in size (0.72 cm), echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.


Liver

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is homogenous echotexture. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

The gall bladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and likely incidental at this time. The cystic and common bile ducts are normal/not visible.

Gastrointestinal

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.36cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is



PATIENT

Mama Agricola

SPECIES

Feline

BREED

DSH

SEX

Spayed Female

AGE

9 years

WEIGHT

12.5 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Pamela Harrigan, RDCS,
Certified Veterinary
Sonographer

HOSPITAL NAME

Norfolk County
Veterinary Service

REFERRING VET

Dr. Jeremy Carignan

INVOICE

10751

DATE

11/13/2025

adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

The visualized areas of duodenum, jejunum and ileum have a uniform diameter with minimal fluid distension. Wall thickness is normal to slightly increased. Bowel loops follow a typical curvilinear path with distinct wall layering, but some areas display a prominent muscularis layer which does not display the typical 1:3 muscularis:mucosa layer ratio. The duodenum measured 0.28 cm in diameter and the jejunum measured 0.3 cm in diameter. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with non-formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

Pancreas

The pancreas is mildly mottled. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

Free Abdomen

Evaluation of the peritoneal cavity did not reveal any evidence of effusion. There is a focal area of inflammation near the ileocecal junction with reactive mesentery and large, hypoechoic, rounded lymph nodes. In this area lymph nodes measure at 0.67 cm, and 0.44 cm in diameter. A smaller lymph node near the ileocecal junction measures 0.24 cm. The omentum is of normal uniform echogenicity.

ULTRASONOGRAPHIC FINDINGS


- Pancreatic changes most consistent with chronic pancreatic remodeling.
- Diffusely “ropey” small intestine with some areas exhibiting a mildly prominent muscularis layer. Findings are most consistent with an inflammatory type enteropathy. Although early neoplastic change cannot be ruled out.
- Cluster of large, hypoechoic lymph nodes and reactive mesentery visualized in the mid right abdomen near the ileocecal junction.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

There is a focus of inflammatory mesentery and large, hypoechoic lymph nodes near the ICJ. If a safe window for sampling is available, consider a fine needle aspirate of a mesenteric lymph node (these may be too small but could be worth trying.) A source of this inflammation is not readily visualized. In general, the small intestine appears somewhat “ropey”, possibly consistent with a mild inflammatory enteropathy, but no focal lesions are observed. Consider the following:

- Consider a novel protein/hydrolyzed protein diet (exclusively at least 4-6 weeks.)
- Consider a GI panel to Texas A&M for evaluation of B12 levels, folate, PLI/TLI etc.. to further evaluate for pancreatic/small intestinal disease.
- Recommend chronic probiotic therapy.
- If not already done, recommend empirical deworming and screening.

If symptoms are persistent, biopsies of the colon and GI tract may eventually be warranted. Prior to this consider repeat imaging looking for the progression of the lymph node enlargement, or the



PATIENT

development of new lesions.

Mama Agricola

SPECIES

Feline

BREED

DSH

SEX

Spayed Female

AGE

9 years

WEIGHT

12.5 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Pamela Harrigan, RDCS,
Certified Veterinary
Sonographer

HOSPITAL NAME

Norfolk County
Veterinary Service

REFERRING VET


Dr. Jeremy Carignan


INVOICE

10751

DATE

11/13/2025





PATIENT

Mama Agricola

SPECIES

Feline

BREED

DSH

SEX

Spayed Female

AGE

9 years

WEIGHT

12.5 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Pamela Harrigan, RDCS,
Certified Veterinary
Sonographer

HOSPITAL NAME

Norfolk County
Veterinary Service

REFERRING VET


Dr. Jeremy Carignan

INVOICE

10751

DATE

11/13/2025



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

info@sonopath.com