



PATIENT

Scooby Clinton

SPECIES

Canine

BREED

Boxer x

SEX

Neutered Male

AGE

13 Years

WEIGHT

82.4 lbs

INTERPRETED BY

Kathleen Sennello DVM,
 MS, Diplomate ACVIM
 (Small Animal Internal
 Medicine)

IMAGING PERFORMED BY

Kathleen Byrnes

HOSPITAL NAME

Shallowford Animal
 Hospital

REFERRING VET

Dr. Eads

INVOICE

71993

DATE

11/20/25

PRESENTING CLINICAL SIGNS

P presented 11/17/25 for ADR and possible seizure, vomiting, abd tight rad report- cardio structures normal size and shape, reduced serosal definition abdomen, wispy appearance ventrally, size of spleen normal however there is an irregular margination with undulating borders to the spleen, The liver is mild to mod enlarged Drained a large amount of fluid- P weighed 77.6# post abdominocentesis P treated with steroids, B12, Cerenia, and fluids P presented 11/20/25 for US- rDVM requests DOUBLE Cavity with Comprehensive echo

Abnormal PE/Chem/CBC/UA Results: TP 4.4, Alb 2.5, ALKP 294, SDMA 16.1, WBC 15.6, Neu 11,388, Monocytes 1,092 T4 0.6 Urinalysis usg 1.009, Protein 2+, Cocci and Rods

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The prostate is normal in size (1.23 cm) and shape for this neutered male dog. The parenchyma is homogenous and the external margins are smooth. The prostatic urethra appears normal with no evidence of irregularity, invasion, mass effect or calculi.

The left kidney has a normal shape and size (6.31 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (6.6 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.


Adrenal Glands

The left adrenal gland is large, measuring 0.89 cm at the cranial pole and 0.86 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is large, measuring 1.4 cm at the cranial pole and 0.90 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

Spleen

The spleen is subjectively normal in size (2.14 cm), echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.



PATIENT

Scooby Clinton

SPECIES

Canine

BREED

Boxer x

SEX

Neutered Male

AGE

13 Years

WEIGHT

82.4 lbs

INTERPRETED BY

Kathleen Sennello DVM,
 MS, Diplomate ACVIM
 (Small Animal Internal
 Medicine)

IMAGING PERFORMED BY

Kathleen Byrnes

HOSPITAL NAME

Shallowford Animal
 Hospital

REFERRING VET

Dr. Eads

INVOICE

71993

DATE

11/20/25

Liver

The liver is large in size, and normal in echogenicity with smooth peripheral margins. The parenchyma is heterogenous in echotexture with subtle, indistinct focal mottling. The visible portions of the vasculature and biliary tract appear normal. There are numerous ill-defined hypoechoic nodules visualized within the parenchyma. Examples of two nodules measure 2.12 cm x 2.99 cm and 2.37 cm in diameter.

The gall bladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and likely incidental at this time. The cystic and common bile ducts are normal/not visible.

Gastrointestinal

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. Jejunum wall measures 0.36 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

Pancreas

The area of the pancreas is normal and isoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

Free Abdomen

There is a large amount of anechoic free fluid. There are occasional prominent mesenteric lymph nodes, an example measures 1.11 cm in diameter. The omentum is mildly diffusely hyperechoic.

ULTRASONOGRAPHIC FINDINGS

- Bilateral adrenomegaly – The bilateral adrenomegaly could be consistent with bilateral hyperplasia (e.g., secondary to pituitary-dependent hyperadrenocorticism), bilateral infiltrative neoplasia, inflammatory adrenal disease, other. Correlation with clinical findings is recommended.
- Heterogeneous liver with ill-defined hypoechoic nodules – The diffuse hepatic changes are non-specific and could be consistent with vacuolar hepatopathy, nodular hyperplasia, inflammatory/immune-mediated disease, fibrosis, extramedullary hematopoiesis, toxic hepatopathy (e.g., copper), infiltrative neoplasia (less likely) or other hepatopathy. The nodules observed trend toward a more benign process but underlying neoplasia cannot be ruled out.
- Occasional prominent mesenteric lymph nodes – These have the appearance most consistent with reactive lymph nodes. Early neoplastic lymph nodes cannot be ruled out.



PATIENT

Scooby Clinton

- Large volume free abdominal fluid.

SPECIES

Canine

BREED

Boxer x

SEX

Neutered Male

AGE

13 Years

WEIGHT

82.4 lbs

INTERPRETED BY

Kathleen Sennello DVM,
 MS, Diplomate ACVIM
 (Small Animal Internal
 Medicine)

IMAGING PERFORMED BY

Kathleen Byrnes

HOSPITAL NAME

Shallowford Animal
 Hospital

REFERRING VET

Dr. Eads

INVOICE


71993

DATE

11/20/25

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

Recommend fluid sampling for fluid analysis and cytology, as well as a cardiac ultrasound to assess for underlying cardiac disease. No large focal mass lesions were observed on today's exam. The liver is somewhat heterogeneous. The significance of this is uncertain. Correlate with clinical signs, current lab work, etc.





PATIENT

Scooby Clinton

SPECIES

Canine

BREED

Boxer x

SEX

Neutered Male

AGE

13 Years

WEIGHT

82.4 lbs

INTERPRETED BY

Kathleen Sennello DVM,
 MS, Diplomate ACVIM
 (Small Animal Internal
 Medicine)

IMAGING PERFORMED BY

Kathleen Byrnes

HOSPITAL NAME

Shallowford Animal
 Hospital

REFERRING VET

Dr. Eads

INVOICE

71993

DATE

11/20/25



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

info@sonopath.com