



PATIENT

Moana Nevarez

SPECIES

Canine

BREED

Miniature Schnauzer

SEX

Spayed Female

AGE

8 Years

WEIGHT

13.4 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Gabriel Ferrer, DVM

HOSPITAL NAME

Pulse: Pet Ultrasound

REFERRING VET

Dr. Jose Vientos

INVOICE

72529

DATE

12/11/25

PRESENTING CLINICAL SIGNS

Pt presented as a referral for an abdominal ultrasound to evaluate hx of intermittent episodes of vomiting and bloody stool. Symptoms stop with treatment for a while but eventually come back. Pt has history of pancreatitis.

Abnormal PE/Chem/CBC/UA Results: CBC: WNL CHEM: WNL CPL: WNL

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, or masses. In the dependent portion of the urinary bladder there is a small amount of dependent sandy debris/small calculi. Some measure up to 0.18 cm in diameter.

The left kidney has a normal shape and size (4.29 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (4.21 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

Adrenal Glands

The left adrenal gland is normal/borderline "flat" measuring 0.35 cm at the cranial pole and 0.41 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is normal/borderline "flat" measuring 0.36 cm at the cranial pole and 0.32 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

Spleen

The spleen is subjectively normal in size (1.25 cm), echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

Liver

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is homogenous echotexture. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.



PATIENT

Moana Nevarez

The gall bladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. There is a moderate amount of non-organized echogenic debris. The cystic and common bile ducts are normal/not visible.

SPECIES

Canine

Gastrointestinal

The stomach contains moderate fluid and gas. It measures at a normal thickness of 0.28 cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. Gas in some areas creates an artifact, limiting evaluation of some areas of the stomach.

BREED

Miniature Schnauzer

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal to mild fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. Duodenum wall measures 0.44 cm. Jejunum wall measures 0.36 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

SEX

Spayed Female

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

AGE

8 Years

Pancreas

The pancreas is normal and isoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

WEIGHT

13.4 lbs

Free Abdomen

Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

ULTRASONOGRAPHIC FINDINGS

IMAGING PERFORMED BY

Gabriel Ferrer, DVM

- Dependent mineralized debris/small stones visualized in the urinary bladder – Recommend urinalysis, culture +/- radiographs.
- Borderline “flat” adrenal glands – Recommend a baseline cortisol.
- Fluid and gas distended stomach – Findings are most concerning for delayed gastric emptying or possibly an unseen partial outflow tract obstruction.

HOSPITAL NAME

Pulse: Pet Ultrasound

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The majority of today’s exam appears relatively normal. The most significant finding is the significantly fluid and gas distended stomach despite an appropriate fast. The proximal duodenum looks normal, and the majority of the visualized stomach appears normal. The exact junction is not definitively visualized. An outflow tract obstruction is unlikely but cannot be definitively ruled out.

REFERRING VET

Dr. Jose Vientos

INVOICE

72529

DATE

12/11/25

Further evaluation could include an upper GI endoscopy combined with colonoscopy to evaluate the stomach, outflow tract obstruction, and obtain biopsies of the proximal GI tract as well as to consider biopsies of the distal GI tract due to the hemorrhagic diarrhea reported. Additional considerations could include the following:



PATIENT

Moana Nevarez

SPECIES

Canine

BREED

Miniature Schnauzer

SEX

Spayed Female

AGE

8 Years

WEIGHT

13.4 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Gabriel Ferrer, DVM

HOSPITAL NAME

Pulse: Pet Ultrasound

REFERRING VET

Dr. Jose Vientos


INVOICE

72529

DATE

12/11/25

- Consider a novel protein/hydrolyzed protein diet (exclusively at least 4-6 weeks)
- Consider a GI panel to Texas A&M for evaluation of B12 levels, folate, PLI/TLI etc.. to further evaluate for pancreatic/small intestinal disease.
- Recommend parasite screening and empirical deworming.
- Recommend a baseline cortisol to screen for Addison's.
- Consider an infectious disease panel for diarrhea.





PATIENT

Moana Nevarez

SPECIES

Canine

BREED

Miniature Schnauzer

SEX

Spayed Female

AGE

8 Years

WEIGHT

13.4 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Gabriel Ferrer, DVM

HOSPITAL NAME

Pulse: Pet Ultrasound

REFERRING VET


Dr. Jose Vientos

INVOICE

72529

DATE

12/11/25



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

info@sonopath.com