



DATE PRESENTING CLINICAL SIGNS

12/21/22 Geriatric screen, has recently had a decreased appetite in the mornings which is unlike him. Also had a syncopal episode during jugular venipuncture 11/2022. Splenectomy - benign cavitated mass 4/2020
PATIENT Pulmonary bullae on x-rays - 4/2020. CKD diagnosed recently on lab work.

Jax Murray Current Medications: Carprofen PRN, Apoquel PRN.
Lab Results: Azotemia, NR anemia.

SPECIES Date of Previous IntraPet Ultrasound: No previous.
Sedation: Not required to complete full diagnostic ultrasound.
Stat Report: Not requested.
Canine

BREED ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Golden Retriever **Urinary System**

SEX The urinary bladder is moderately distended with anechoic urine. The Bladder wall is diffusely mildly thickened (0.52 cm), and the mucosa is mildly irregular. The trigone, ureteral papillae, and visible urethra (to a depth of 2cm) appear normal with no evidence of severe mucosal irregularities, masses or cystic calculi. Findings are most consistent with bacterial cystitis or lack of urine distension. Recommend urinalysis and culture.

Neutered Male

AGE The prostate is not clearly seen.

5/4/08

WEIGHT The left kidney has a normal shape and size (6.19 cm) with pyelectasia at 0.50 cm. Overall echogenicity is slightly hyperechoic with poor corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of nephroliths, infarcts or hydroureter. Renal vasculature is normal.

64 Pounds

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

The right kidney has a normal shape and size (6.32 cm) with pyelectasia at 0.39 cm. Overall echogenicity is slightly hyperechoic with poor corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of nephroliths, infarcts or hydroureter. Renal vasculature is normal.

IMAGING PERFORMED BY

Stephanie Warga
RDMS, RVT

Adrenal Glands

The left adrenal gland is normal in size measuring 0.96 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

HOSPITAL NAME

Stay Pet Veterinary

The right adrenal gland is normal in size measuring 0.70 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

REFERRING VET

Dr. Klimovitz

Spleen

The spleen is surgically absent, splenectomy performed 4/2020.

Liver

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is heterogenous in echotexture with subtle, indistinct focal mottling. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

INVOICE

43611

The gall bladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. There is a moderate amount of non-organized echogenic debris. The proximal bile duct appears somewhat prominent and dilated at 0.91 cm.

Gastrointestinal

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. Duodenum wall measures 0.46 cm. Jejunum wall measures 0.31 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

Pancreas

The pancreas is normal and isoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

Free Abdomen

Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

Other

A brief view of the heart was submitted. No significant pericardial effusion was seen.

PRIMARY FINDINGS

- Mildly thickened/irregular urinary bladder wall – The bladder mucosal changes could be consistent with cystitis or artifactual due to lack of adequate luminal distension. Bladder neoplasia cannot be ruled out but is considered unlikely in this patient.
- Decreased corticomedullary distinction in both kidneys with bilateral pyelectasia – Mild loss of corticomedullary distinction in both kidneys could be consistent with chronic degenerative disease or interstitial nephrosis. Pyelectasia of the kidney(s) could be consistent with pyelonephritis, chronic renal disease, secondary to PU/PD or fluid therapy (if applicable), other.
- Heterogeneous liver – The diffuse hepatic changes are non-specific and could be consistent with vacuolar hepatopathy, nodular hyperplasia, inflammatory/immune-mediated disease, fibrosis, extramedullary hematopoiesis, toxic hepatopathy (e.g., copper), infiltrative neoplasia (less likely) or other hepatopathy.
- Moderate gallbladder debris with a mildly dilated proximal bile duct – The significance of the aggregated gallbladder debris is unclear. This could represent an early mucocele, cholestasis, or may be secondary to fasting but seems unlikely to be causing a current issue. Recommend continued monitoring.

SECONDARY FINDINGS

- Surgically absent spleen – Splenectomy performed 4/2020 for a benign mass lesion.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS


No overt mass lesions are observed, or evidence of metastatic disease. The liver appears somewhat heterogeneous, but this is a non-specific finding and could be associated with age related remodeling.


The changes in the kidneys are consistent with chronic age related progressive renal disease. Bilateral pyelectasia could be secondary to PU/PD, but a urinalysis and culture is recommended to rule out pyelonephritis. Additionally, a blood pressure should be evaluated to establish baseline and look for evidence of hypertension.

The urinary bladder wall appears slightly thickened and irregular. Recommend a urinalysis and culture (as recommended above).

There is a moderate amount of debris in the gallbladder, and the proximal bile duct appears slightly prominent. With no evidence of significant liver enzyme elevations, this can continue to be monitored.

Keep in mind that NSAID therapy may be contraindicated in this patient, as it could increase risk for GI ulceration and acute on chronic kidney disease. You could consider non-steroidal pain medications, although sometimes this becomes a quality of life issue, and comfort is prioritized.





The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)
kathleen.sennello@sonopath.com