



PATIENT

Butters Collins

SPECIES

Feline

BREED

British Shorthair

SEX

Intact Male

AGE

4 Months

WEIGHT

1.87 kg

INTERPRETED BY

Tam Mengine, DVM,
DABVP (canine/feline
practice)

IMAGING PERFORMED BY

Dr. Sarah Barthelemy

HOSPITAL NAME

Fish Creek Pet Hospital

REFERRING VET

Dr. Johnson

INVOICE

72796

DATE

12/28/25

PRESENTING CLINICAL SIGNS

Patient presented yesterday for lethargy, inappetence, not wanting to be held, and some wheezing/sneezing. Was mildly febrile and decision was for outpatient treatment for possible upper Resp infection as litter mate has similar signs. Metacam injection given and 5x overdose was administered by mistake. Patient then hospitalized on IVF, etc and labs showed elevated globulins, elevated ALT and tbili, pleural effusion identified and centesis revealed yellow tinged fluid with protein. FeLV and FIV negative. FIP PCR pending

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine, and no luminal sediment is present. The ureteral papillae, trigone and pelvic urethra (visible to 2.0 cm) are of normal appearance, and the ureters are not visible (normal). No masses, calculi or mucosal irregularities are noted.

The kidneys are subjectively enlarged relative to patient size, with appropriate shape and corticomedullary differentiation with a normal 1:3 cortex to medulla ratio. There are subtle striations noted at the corticomedullary junction. The proximal ureters are not visible (normal). Left kidney measures 3.6 cm. Right measures 3.9 cm.

Adrenal Glands

The adrenal glands are both identified in their normal locations. They are normal in size and shape with appropriate parenchymal echogenicity and normal phrenic vasculature. Left measures 3.7 mm. Right measures 3.3 mm.

Spleen

The spleen is of appropriate size (6.3 mm) and has a normal, homogenous parenchyma with a smooth, continuous capsular surface. The splenic vasculature is normal with no evidence of congestion or thrombosis, and blood flow through the splenic hilus appears normal. Thickness at the splenic hilus is normal.

Liver

The liver is of appropriate size and shape, with a diffusely hypoechoic parenchyma and increased portal markings. The portal and hepatic vasculature are of normal size and appearance with no evidence of congestion or thrombosis.

The gallbladder is moderately distended with anechoic contents. The wall was thin and continuous with no focal lesions. The cystic and common bile ducts are normal / not visible.

Gastrointestinal

The stomach is empty. The gastric wall is subjectively normal in thickness, and exhibits appropriate wall layering, but cannot be accurately measured due to normal deviations of the rugal folds. The pylorus is of normal appearance.

The visualized portions of the duodenum, jejunum, and ileum are of normal thickness with intact wall layering that exhibits the appropriate 1:3 muscularis to mucosa ratio. Intestinal motility appears normal. Duodenum wall measures 1.5 mm. Jejunum wall measures 2.5 mm.



PATIENT

Butters Collins

SPECIES

Feline

BREED

British Shorthair

SEX

Intact Male

AGE

4 Months

WEIGHT

1.87 kg

INTERPRETED BY

Tam Mengine, DVM,
DABVP (canine/feline
practice)

**IMAGING
PERFORMED BY**

Dr. Sarah Barthelemy

HOSPITAL NAME

Fish Creek Pet Hospital

REFERRING VET

Dr. Johnson

INVOICE

72796

DATE

12/28/25



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Tam Mengine, DVM, DABVP (canine/feline practice)

info@SonoPath.com