



PATIENT

Nala Tahis Velazquez

SPECIES

Canine

BREED

Toy Poodle

SEX

Spayed Female

AGE

12

WEIGHT

8.3 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Gabriel Ferrer, DVM

HOSPITAL NAME

Pulse: Pet Ultrasound

REFERRING VET

Dr. Hector Perez

INVOICE

72321

DATE

12/4/25

PRESENTING CLINICAL SIGNS

Pt presented as a referral for an abdominal ultrasound to evaluate loss of appetite that started 6 days ago. Increased water intake and urine (clear, now yellow). Otherwise, pt is BAR and no other clinical signs. Was switched from Science Diet to Royal Canin Gastrointestinal canned food 1-2 days ago.

Abnormal PE/Chem/CBC/UA Results: Bloodwork and radiographs attached as supporting documents.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The left kidney has a normal shape and size (3.26 cm) with mild pyelectasia at 0.13 cm. Overall echogenicity is slightly hyperechoic with poor corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (3.75 cm). Overall echogenicity is slightly hyperechoic with poor corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

Adrenal Glands

The left adrenal gland is normal in size measuring 0.40 cm at the cranial pole and 0.46 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is normal in size measuring 0.59 cm at the cranial pole and 0.49 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

Spleen

The spleen is subjectively normal in size (1.03 cm in width at the level of the hilus), echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. There is a small hypoechoic nodule in the parenchyma measuring 0.22 cm x 0.27 cm.

Liver

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is homogenous echotexture. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.



PATIENT	The gall bladder lumen is significantly distended. Some areas of the wall appear mildly thickened with adherent debris. There is a large amount of primarily non-organized echogenic debris. There is no evidence of bile duct dilation.
Nala Tahis Velazquez	
SPECIES	<i>Gastrointestinal</i>
Canine	The stomach contains mild fluid/shadowing ingesta. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.
BREED	
Toy Poodle	The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. Duodenum wall measures 0.38 cm. Jejunum wall measures 0.31 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.
SEX	
Spayed Female	
AGE	The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.
12	
WEIGHT	<i>Pancreas</i>
8.3 lbs	The pancreas is visible/mildly mottled in the right limb. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.
INTERPRETED BY	<i>Free Abdomen</i>
Kathleen Sennello DVM, MS, Diplomate ACVIM (Small Animal Internal Medicine)	Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.
IMAGING PERFORMED BY	ULTRASONOGRAPHIC FINDINGS
Gabriel Ferrer, DVM	<ul style="list-style-type: none"> • Mild age related changes visualized associated with both kidneys. • Small, hypoechoic nodule in the spleen – There is a non-cavitated, hypoechoic splenic nodule visualized. Differentials include lymphoid hyperplasia, extramedullary hematopoiesis, infiltrative neoplasia, inflammation, other. Cytology or histopathology would be necessary to get a definitive diagnosis. • Pancreatic changes most consistent with chronic pancreatic remodeling. Mild chronic pancreatitis is possible. • Distended gallbladder with a large amount of disorganized luminal debris – A large amount of debris is evident in the gall bladder with no evidence of a mucocele or associated inflammation at this time. This could represent an early mucocele or cholestasis, with minimal evidence of associated inflammation at this time. Continued monitoring of labwork and ultrasound are warranted for progression of this lesion. Ursodiol therapy could be considered.
HOSPITAL NAME	
Pulse: Pet Ultrasound	
REFERRING VET	
Dr. Hector Perez	
INVOICE	
72321	
DATE	
12/4/25	
	<u>INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS</u>
	The changes observed on today's scan are relatively mild. There were mild age related changes



PATIENT

Nala Tahis Velazquez

SPECIES

Canine

BREED

Toy Poodle

SEX

Spayed Female

AGE

12

WEIGHT

8.3 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Gabriel Ferrer, DVM

HOSPITAL NAME

Pulse: Pet Ultrasound

REFERRING VET

Dr. Hector Perez

INVOICE

72321

DATE


12/4/25

visualized associated with the kidneys. Recommend correlating full lab work including electrolytes with urinalysis to better assess the azotemia reported. Either way, supportive fluids are recommended, either as diuresis and/or to rehydrate. Additionally consider a urine culture/urine protein to creatinine ratio and blood pressure evaluation if clinically appropriate.

There is a very small nodule visualized associated with the spleen. I suspect this is too small to easily sample. Recommend continued monitoring with ultrasound.

No evidence of active pancreatitis is present, but the pancreas is visible, and unseen pancreatic inflammation is possible. Consider empirical treatment for pancreatitis and gastroenteritis. If anorexia is persistent, a feeding tube may need to be considered.

There is a large amount of debris in the gallbladder but no evidence of surrounding inflammation or wall thickening. Consider chronic Ursodiol therapy and continued monitoring of lab work and the gallbladder with ultrasound.





PATIENT

Nala Tahis Velazquez

SPECIES

Canine

BREED

Toy Poodle

SEX

Spayed Female

AGE

12

WEIGHT

8.3 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Gabriel Ferrer, DVM

HOSPITAL NAME

Pulse: Pet Ultrasound

REFERRING VET

Dr. Hector Perez

INVOICE

72321

DATE

12/4/25



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

info@sonopath.com