



**PATIENT**

Mia Pevny

**PRESENTING CLINICAL SIGNS**

presented for vomiting after drinking large amounts of water, PU/PD gained ~3.5 lbs increased UCCR and ALT

**SPECIES**

Canine

Abnormal PE/Chem/CBC/UA Results: 11/18/22- LYM: 0.8L, NEU%: 83.7H, LYM%: 9.2L, HCT: 56.2H, ALP: 326H, Potassium: 3.7L 11/29/22- UCCR: 37 12/3- LDDS Pre: 3.9 4hrs post: 2.5 8hrs post: 4.9

**BREED**

Terrier X

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**Urinary System**

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

**SEX**

Spayed Female

The left kidney has a normal shape and size (6.64 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

**AGE**

9 Years 10 Months

The right kidney has a normal shape and size (6.9 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

**WEIGHT**

51.6 Pounds

**Adrenal Glands**

The left adrenal gland is normal in size measuring 0.61 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

The right adrenal gland is normal in size measuring 0.50 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

**IMAGING PERFORMED BY**

Dr. Striano-Kaplan

**Spleen**

The spleen is subjectively normal in size, echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

**HOSPITAL NAME**

Ramsey Vet Hospital

**Liver**

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is homogenous echotexture. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

**REFERRING VET**

Dr. Striano-Kaplan

The gallbladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and primarily anechoic. The cystic and common bile ducts are normal/not visible.

**INVOICE**

43240

**DATE**

12/6/22



**PATIENT**

***Gastrointestinal***

Mia Pevny

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

**SPECIES**

Canine

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. Jejunum wall measures 0.36 cm.

**BREED**

Terrier X

Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

**SEX**

Spayed Female

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

**AGE**

9 Years 10 Months

***Pancreas***

The pancreas is normal and isoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

**WEIGHT**

51.6 Pounds

***Free Abdomen***

Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

**ULTRASONOGRAPHIC FINDINGS**

- No significant ultrasonographic lesions observed

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

Today's scan appears relatively normal for a 10 year old mixed breed dog. No mass lesions were observed. The adrenals appear relatively normal in size. Additionally, no lesions associated with vomiting were observed. Unfortunately, there are many causes for PU/PD that cannot be diagnosed by ultrasound alone. Cushing's disease is a possibility here, but slightly less likely, given normal adrenals. Correlate with clinical signs. Additionally, you could consider an ACTH stimulation test, as this test would have fewer false positives, and consider the possibility of non-adrenal illness.

**IMAGING PERFORMED BY**

Dr. Striano-Kaplan

An obvious lesion responsible for the reported increase in thirst and urination was not visualized. Some issues such as early renal disease, Cushing's disease, behavioral, neurologic, dietary, electrolyte disturbances etc.. are not able to be diagnosed with ultrasound alone. These can be challenging cases.

**HOSPITAL NAME**

Ramsey Vet Hospital

The top 10 differentials can be ruled in/out with routine bloodwork, urinalysis and culture, several more can be evaluated with a good history and imaging. Unfortunately, as you work your way down the list the differentials become harder to definitively diagnose. This is the differential list I start with.

**REFERRING VET**

Dr. Striano-Kaplan

- Diabetes Mellitus
- Chronic Renal Disease/Renal Failure (can present pre-azotemic, especially in dogs, but expect the BUN & creatinine not to be at the low end of the reference range)
- Hypercalcemia
- Urinary tract infection
- Iatrogenic Disease due to medications (diuretics, phenobarbital, KBr; diets either high in salt [such as S/D] or very low in protein (such as U/D))
- Hyperthyroidism
- Hypokalemia

**INVOICE**

43240

**DATE**

12/6/22



**PATIENT**

Mia Pevny

**SPECIES**

Canine

**BREED**

Terrier X

**SEX**

Spayed Female

**AGE**

9 Years 10 Months

**WEIGHT**

51.6 Pounds

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

**IMAGING  
PERFORMED BY**

Dr. Striano-Kaplan

**HOSPITAL NAME**

Ramsey Vet Hospital

**REFERRING VET**

Dr. Striano-Kaplan


**INVOICE**

43240

**DATE**

12/6/22

- Liver Disease (hepatic encephalopathy may be a mixed primary PU and PD)
- Pyelonephritis
- Polycythemia
- Renal Tubular Diseases (glycosuria or Fanconi & Fanconi-like syndromes or RTA)
- Hyperadrenocorticism (may be a mixed primary PU and PD)
- Hypoadrenocorticism (either Addison's or hypocortisolism)
- Paraneoplastic Syndromes (particularly splenic hemangiosarcoma?)
- Pericardial Effusion
- Pyometra (including stump pyometra in spayed dogs)
- Chronic Partial Urinary Obstruction or Post-Obstructive Diuresis
- Pheochromocytoma
- Psychogenic Polydipsia (as in a true behavior disorder with a compulsive element)
- Primary Non-Medical Polydipsia (aka "I drink a lot because I like it or I engage in activities that promote it, but that doesn't mean I'm sick")
- Primary Nephrogenic Diabetes Insipidus (Congenital Nephrogenic Diabetes Insipidus, other diseases that cause primary PU other than Congenital Diabetes Insipidus would be considered Acquired Nephrogenic Diabetes Insipidus)
- Atypical Cushing's and SARDS
- Central Diabetes Insipidus





**PATIENT**

Mia Pevny

**SPECIES**

Canine

**BREED**

Terrier X

**SEX**

Spayed Female

**AGE**

9 Years 10 Months

**WEIGHT**

51.6 Pounds

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

**IMAGING  
PERFORMED BY**

Dr. Striano-Kaplan

**HOSPITAL NAME**

Ramsey Vet Hospital

**REFERRING VET**


Dr. Striano-Kaplan

**INVOICE**

43240

**DATE**

12/6/22



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

kathleen.sennello@sonopath.com