



PATIENT

Casper Cosentino

SPECIES

Feline

BREED

Himalayan

SEX

Neutered Male

AGE

8 Years

WEIGHT

5.5 Pounds

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Kelly Vazquez

HOSPITAL NAME

Companion Vet
Hospital of Wayne

REFERRING VET

Dr. Ben Spitz

INVOICE

44691

DATE

2/1/23

PRESENTING CLINICAL SIGNS

Patient presents for concerns of possible neoplasia. Patient has a history of poorly controlled pemphigus and skin disease; has a heart murmur. Current meds: Gentocin spray, Miconahex + Triz® Mousse, medrol, chlorambucil, Proviavle-Forte®, and gabapentin.

Abnormal PE/Chem/CBC/UA Results: No current blood work.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The left kidney has a normal shape and size (4.04 cm). Overall echogenicity is slightly hyperechoic with poor corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (3.63 cm). Overall echogenicity is slightly hyperechoic with poor corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

Adrenal Glands

The left adrenal gland is normal in size measuring 0.50 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is normal in size measuring 0.48 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

Spleen

The spleen is subjectively normal in size (0.74 cm). It is mottled and hypoechoic, the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

Liver

The liver is large in size, and normal in echogenicity with smooth peripheral margins. The parenchyma is heterogenous in echotexture with subtle, indistinct focal mottling. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

The gallbladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and primarily anechoic. The cystic and common bile ducts are normal/not visible.



PATIENT

Casper Cosentino

SPECIES

Feline

BREED

Himalayan

SEX

Neutered Male

AGE

8 Years

WEIGHT

5.5 Pounds

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Kelly Vazquez

HOSPITAL NAME

Companion Vet
Hospital of Wayne

REFERRING VET

Dr. Ben Spitz

INVOICE

44691

DATE

2/1/23

Gastrointestinal

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.36cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

The visualized areas of duodenum, jejunum and ileum have a uniform diameter with minimal fluid distension. Wall thickness is normal to slightly increased. Bowel loops follow a typical curvilinear path with distinct wall layering, but some areas display a prominent muscularis layer which does not display the typical 1:3 muscularis:mucosa layer ratio. Duodenum wall measures 0.33 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

Pancreas

There is a large, slightly hypoechoic, mottled mass effect visualized in the abdomen running from the caudal aspect of the stomach cranially down to the gastroduodenal flexure. No normal pancreas is visualized in this region, and I suspect this is a pancreatic mass effect. Adjacent to the duodenum, there is a small area of more normal hypoechoic pancreas, which comes into contact with this abnormal tissue. This mass effect measures approximately 1.72 cm x >7.0 cm in length.

Free Abdomen

Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

ULTRASONOGRAPHIC FINDINGS

- Decreased corticomedullary distinction in both kidneys – Mild loss of corticomedullary distinction in both kidneys could be consistent with chronic degenerative disease or interstitial nephrosis.
- Hypoechoic mottled spleen – The diffuse splenic changes are non-specific and could be consistent with lymphoid hyperplasia, extramedullary hematopoiesis, infiltrative neoplasia, inflammation, other. Cytology or histopathology would be necessary to get a definitive diagnosis.
- Large, heterogeneous, hypoechoic mass effect in the region of the left limb of the pancreas – Findings are concerning for a likely pancreatic mass effect. Differentials would include carcinoma, round cell neoplasia, etc.
- Heterogeneous liver – The diffuse hepatic changes are non-specific and could be consistent with vacuolar hepatopathy, nodular hyperplasia, inflammatory/immune-mediated disease, fibrosis, extramedullary hematopoiesis, toxic hepatopathy (e.g., copper), infiltrative neoplasia (less likely) or other hepatopathy.
- Prominent muscularis layer to the small intestine – The small intestinal wall changes could be consistent with an underlying inflammatory process. These types of changes can sometimes be seen in normal older cats. Correlate with clinical signs.



PATIENT

Casper Cosentino

SPECIES

Feline

BREED

Himalayan

SEX

Neutered Male

AGE

8 Years

WEIGHT

5.5 Pounds

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Kelly Vazquez

HOSPITAL NAME

Companion Vet
Hospital of Wayne

REFERRING VET

Dr. Ben Spitz

INVOICE

44691

DATE


2/1/23

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

There is a very large mixed echogenic hypoechoic mass effect in the cranial abdomen running caudal to the stomach and down into the area of the portal hilus. This could correlate with the skin lesions, causing paraneoplastic lesions. Recommend a fine needle aspirate of the mass effect.

Additionally, the spleen appears somewhat heterogeneous and hypoechoic, and the liver is mildly heterogeneous. If a cytologic diagnosis cannot be obtained based on pancreatic aspirate, consider an aspirate of the spleen.

Recommend three view thoracic radiographs to evaluate for possible concurrent thoracic disease/involvement.





PATIENT

Casper Cosentino

SPECIES

Feline

BREED

Himalayan

SEX

Neutered Male

AGE

8 Years

WEIGHT

5.5 Pounds

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Kelly Vazquez

HOSPITAL NAME

Companion Vet
Hospital of Wayne

REFERRING VET


Dr. Ben Spitz

INVOICE

44691

DATE

2/1/23



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

kathleen.sennello@sonopath.com