



PATIENT

Feli Hermo

SPECIES

Canine

BREED

Mixed

SEX

Neutered Male

AGE

10 Years

WEIGHT

51 lbs

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Meghan Morse, LVT,
CVT

HOSPITAL NAME

Hamburg Veterinary
Clinic

REFERRING VET

Hamburg Veterinary
Clinic

INVOICE

73011

DATE

2/17/26

PRESENTING CLINICAL SIGNS

Diarrhea. Inappetence. Increased Urination. Current meds: Metronidazole

Abnormal PE/Chem/CBC/UA Results: ALT 1155, SAP 1662, WBC 21K.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The prostate is normal in size (0.75 cm) and shape for this neutered male dog. The parenchyma is homogenous and the external margins are smooth. The prostatic urethra appears normal with no evidence of irregularity, invasion, mass effect or calculi.

The left kidney has a normal shape and size (5.77 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (5.68 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

Adrenal Glands

The left adrenal gland is normal in size measuring 0.78 cm at the cranial pole and 0.69 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is normal in size measuring 1.18 cm at the cranial pole and 0.83 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

Spleen

The spleen is subjectively normal in size (2.01 cm), echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

Liver

The liver is large in size and irregular in shape. The parenchyma is heterogenous in echotexture with subtle, indistinct focal mottling. The visible portions of the vasculature and biliary tract appear normal. There is a large, mixed echogenicity, hypoechoic, cavitated mass effect that appears to be arising from the caudal left aspect of the liver measuring >6.07 cm x 4.71 cm. There is a 2nd irregular, solid, hypoechoic mass effect that appears to be arising from a similar area in the liver, measuring approximately 5.58 cm x 5.1 cm.



PATIENT

Feli Hermo

The gall bladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and likely incidental at this time. The cystic and common bile ducts are normal/not visible.

SPECIES

Canine

Gastrointestinal

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

BREED

Mixed

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. Duodenum wall measures 0.47 cm. Jejunum wall measures 0.28 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

SEX

Neutered Male

AGE

10 Years

Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

Pancreas

WEIGHT

51 lbs

The right limb of the pancreas is prominent and mottled compared to the surrounding isoechoic mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

Free Abdomen

There is scant free fluid noted. No lymphadenopathy noted. The omentum is diffusely hyperechoic.

The right auricle and pericardium were visualized and were unremarkable. No obvious pathology is visualized. If cardiac function evaluation is desired a full echocardiogram is warranted.

IMAGING PERFORMED BY

Meghan Morse, LVT,
CVT

ULTRASONOGRAPHIC FINDINGS

- Large, irregular, heterogeneous liver with a large, hypoechoic, mixed echogenicity, cavitated mass lesion, and a 2nd solid mass effect – Findings are concerning for possible neoplastic lesion (hemangiosarcoma, carcinoma, other).
- Pancreatic changes most consistent with chronic pancreatic remodeling +/- chronic pancreatitis.

HOSPITAL NAME

Hamburg Veterinary
Clinic

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

There is a very large, mixed echogenicity, hypoechoic, cavitated mass effect in the left cranial abdomen. This appears to be arising from the liver. There is a 2nd more solid mass effect visualized caudal to this lesion, which appears to be hepatic in origin as well. Strongly recommend a contrast CT scan of the abdomen to confirm the origins and nature of lesions observed for surgical planning. Additionally, recommend 3-view thoracic radiographs, looking for any evidence of metastatic disease or pulmonary disease.

REFERRING VET

Hamburg Veterinary
Clinic

INVOICE

73011

No definitive primary gastrointestinal lesions are observed, although the left limb of the pancreas is somewhat prominent. Consider concurrent empirical treatment for gastroenterocolitis and pancreatitis.

DATE

2/17/26



PATIENT

Feli Hermo

SPECIES

Canine

BREED

Mixed

SEX

Neutered Male

AGE

10 Years

WEIGHT

51 lbs

INTERPRETED BY

Kathleen Sennello DVM,
 MS, Diplomate ACVIM
 (Small Animal Internal
 Medicine)

**IMAGING
 PERFORMED BY**

Meghan Morse, LVT,
 CVT

HOSPITAL NAME

Hamburg Veterinary
 Clinic

REFERRING VET


Hamburg Veterinary
 Clinic

INVOICE

73011

DATE

2/17/26





PATIENT

Feli Hermo

SPECIES

Canine

BREED

Mixed

SEX

Neutered Male

AGE

10 Years

WEIGHT

51 lbs

INTERPRETED BY

Kathleen Sennello DVM,
 MS, Diplomate ACVIM
 (Small Animal Internal
 Medicine)

IMAGING PERFORMED BY

Meghan Morse, LVT,
 CVT

HOSPITAL NAME

Hamburg Veterinary
 Clinic

REFERRING VET


Hamburg Veterinary
 Clinic

INVOICE

73011

DATE

2/17/26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

info@sonopath.com