



PATIENT

Killer Kloc

SPECIES

Feline

BREED

DSH

SEX

Spayed Female

AGE

14 Years 10 Months

WEIGHT

11.8

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Dr. Miranda Fritz

HOSPITAL NAME

Waterbury VH

REFERRING VET

Dr. Miranda Fritz

INVOICE

46224

DATE

3/28/23

PRESENTING CLINICAL SIGNS

P had soft tissue sarcoma on her right hind paw, lateral aspect of her metatarsals. It was surgically removed along with the associated 5th digit May 2022. In January 2023 there was regrowth in that region. Biopsy done and confirmed recurrence of the sarcoma. Today we are staging for limb amputation. P sensitive and lame on RH leg but otherwise doing well at home. No recent v/d/c/s, no pu/pd. Previously diagnosed hyperthyroidism (I-131tx Oct 2021) and secondary hyperparathyroidism based on MSU hypercalcemia panel

Abnormal PE/Chem/CBC/UA Results: PE - RH lateral paw semifirm 1cm movable nodule, no osteolysis on x-ray CBC - wnl Chem - total calcium 11.8 UA - USG 1.021, trace protein, all else wnl PTH 8.0 (H) iCa 1.19 (normal) PTH-rp 0.0 (normal) Chest x-rays - wnl

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The left kidney has a normal shape and size (3.29 cm) with mild mineralization at the level of the pelvis. Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (3.68 cm) with mild mineralization at the level of the pelvis. Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, infarcts or hydroureter. Renal vasculature is normal.

Adrenal Glands

The region of left adrenal (Cranial to left renal artery) is unremarkable but the adrenal is not distinctly visualized. No evidence of a mass effect is visualized.

The right adrenal gland is normal in size measuring 0.48 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

Spleen

The spleen is subjectively normal in size (0.98 cm), echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

Liver

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is homogenous echotexture. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

The gallbladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and primarily anechoic. The cystic and common bile ducts are normal/not visible.



PATIENT

Killer Kloc

SPECIES

Feline

BREED

DSH

SEX

Spayed Female

AGE

14 Years 10 Months

WEIGHT

11.8

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Dr. Miranda Fritz

HOSPITAL NAME

Waterbury VH

REFERRING VET

Dr. Miranda Fritz

INVOICE

46224

DATE

3/28/23

Gastrointestinal

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.36cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. The duodenum measured as normal (between 0.13-0.38cm in wall thickness) and the jejunum measured as normal (between 0.15-0.36cm.) Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

Pancreas

The area of the pancreas is normal and isoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

Free Abdomen


Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

ULTRASONOGRAPHIC FINDINGS

- No significant ultrasonographic lesions observed

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

There were no lesions visualized consistent with a metastatic process. There is mild mineralization in both kidneys, which is likely incidental at this time.





PATIENT

Killer Kloc

SPECIES

Feline

BREED

DSH

SEX

Spayed Female

AGE

14 Years 10 Months

WEIGHT

11.8

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Dr. Miranda Fritz

HOSPITAL NAME

Waterbury VH

REFERRING VET


Dr. Miranda Fritz

INVOICE

46224

DATE

3/28/23



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

kathleen.sennello@sonopath.com