

PATIENT

Riggs Jefferies

SPECIES

Canine

BREED

Shiloh Shepherd

SEX

Neutered Male

AGE

2 Years

WEIGHT

56 kg

INTERPRETED BY

Kathleen Sennello DVM,
 MS, Diplomate ACVIM
 (Small Animal Internal
 Medicine)

IMAGING PERFORMED BY

Amanda Stewart

HOSPITAL NAME

Graham Animal
 Hospital

REFERRING VET

Dr. Seager

INVOICE

74079

DATE

3/31/26

PRESENTING CLINICAL SIGNS

Presented for Neuter, clinically normal, wellness BW showed increase creatine, repeated after surgery and had increased. UPC normal. UA normal other than mild increase in WBC. Brief ultrasound showed possible abnormal R kidney

Current Medications: Aventi Kidney 2.5 lg scoops BID, amoxi-clav 500/125 1.25 BID

Abnormal PE/Chem/CBC/UA Results: Initial Creat 170Umol/L 3/13/26 2nd Creat 196 umol/L 3/26/26
 Primary Question to Be Answered in This Exam Cause of renal disease acquired vs congenital and prognosis.

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The visualized areas of prostate and surrounding tissue appear normal. Unfortunately, the prostate is not fully visualized likely due to its intrapelvic location. Correlate with rectal exam findings.

The left kidney has a normal shape and size (8.19 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (8.35 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.


Adrenal Glands

The left adrenal gland is normal/borderline "flat" measuring 0.38 cm at the cranial pole and 0.44 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is normal in size measuring 1.27 cm at the cranial pole and 0.50 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

Spleen

The spleen is normal/borderline large (4.12 cm), echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.



PATIENT

Riggs Jefferies

SPECIES

Canine

BREED

Shiloh Shepherd

SEX

Neutered Male

AGE

2 Years

WEIGHT

56 kg

INTERPRETED BY

Kathleen Sennello DVM,
 MS, Diplomate ACVIM
 (Small Animal Internal
 Medicine)

IMAGING PERFORMED BY

Amanda Stewart

HOSPITAL NAME

Graham Animal
 Hospital

REFERRING VET

Dr. Seager

INVOICE

74079

DATE

3/31/26

Liver

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is homogenous echotexture. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

The gall bladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and likely incidental at this time. The cystic and common bile ducts are normal/not visible.

Gastrointestinal

The visualized area of stomach appears to contain minimal luminal contents. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. The duodenum measured as normal (between 0.3-0.5cm in wall thickness) and the jejunum measured as normal (between 0.2-0.47cm.) Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

Pancreas

The area of the pancreas is normal and isoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

Free Abdomen


Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

ULTRASONOGRAPHIC FINDINGS

- Prominent/borderline large spleen – This could be normal for such a large dog. No focal lesions are visualized. Other less likely differentials could include congestion, splenitis, lymphoid hyperplasia, or infiltrative neoplasia. A fine needle aspirate could be considered if this is a strong concern.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

No significant ultrasonographic lesions are visualized associated with the kidneys. Unfortunately, there can still be a significant nephropathy despite relatively normal appearance of the kidneys. The significance of the creatinine elevation needs to be evaluated in light of urine concentrating ability (urine specific gravity), hydration status, etc. An SDMA level could be helpful. Additionally recommend a baseline cortisol to screen for Addison's.



PATIENT

Riggs Jefferies

SPECIES

Canine

BREED

Shiloh Shepherd

SEX

Neutered Male

AGE

2 Years

WEIGHT

56 kg

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Amanda Stewart

HOSPITAL NAME

Graham Animal
Hospital

REFERRING VET


Dr. Seager


INVOICE

74079

DATE

3/31/26





PATIENT

Riggs Jefferies

SPECIES

Canine

BREED

Shiloh Shepherd

SEX

Neutered Male

AGE

2 Years

WEIGHT

56 kg

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Amanda Stewart

HOSPITAL NAME

Graham Animal
Hospital

REFERRING VET

Dr. Seager

INVOICE

74079

DATE

3/31/26

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

info@sonopath.com