



**PATIENT**

Scarlett Walls

**SPECIES**

Canine

**BREED**

Shar Pei

**SEX**

Spayed Female

**AGE**

8 Years

**WEIGHT**

44 Pounds

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

**IMAGING  
PERFORMED BY**

Sorbo

**HOSPITAL NAME**

Mill Brook AC

**REFERRING VET**

Dr. Jeffers

**INVOICE**

45769

**DATE**

3/8/23

**PRESENTING CLINICAL SIGNS**

Breeding bitch until Dec 2021 when had OVH as part of pyometra Sx. Creatinine at that time was 0.7mg/dL. Now: PUPD and accidents in the house when stressed only. On renal diet. Abnormal PE/Chem/CBC/UA Results: Current labs: creatinine 2.5mg/dL, SDMA 24.4, USG 1.008.

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**Urinary System**

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The left kidney has a normal shape and size (4.92 cm). Overall echogenicity is slightly hyperechoic with poor corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (5.44 cm). Overall echogenicity is slightly hyperechoic with poor corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

**Adrenal Glands**

The left adrenal gland is normal in size measuring 0.52 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is normal in size measuring 0.58 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

**Spleen**

The spleen is subjectively normal in size, echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

**Liver**

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is homogenous echotexture. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

The gallbladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and primarily anechoic. The cystic and common bile ducts are normal/not visible.

**Gastrointestinal**

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.



**PATIENT**

Scarlett Walls

**SPECIES**

Canine

**BREED**

Shar Pei

**SEX**

Spayed Female

**AGE**

8 Years

**WEIGHT**

44 Pounds

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

**IMAGING PERFORMED BY**

Sorbo

**HOSPITAL NAME**

Mill Brook AC

**REFERRING VET**

Dr. Jeffers

**INVOICE**

45769

**DATE**

3/8/23

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. The duodenum measured as normal (between 0.3-0.5cm in wall thickness) and the jejunum measured as normal (between 0.2-0.47cm.) Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

**Pancreas**

The left limb of the pancreas is prominent and hypoechoic as compared to the surrounding isoechoic mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

**Free Abdomen**

Evaluation of the peritoneal cavity did not reveal any evidence of effusion. There are visible/mildly enlarged mesenteric lymph nodes measuring 0.45 cm and 0.86 cm. The omentum is of normal echogenicity.

**ULTRASONOGRAPHIC FINDINGS**

- Decreased corticomedullary distinction in both kidneys – Mild loss of corticomedullary distinction in both kidneys could be consistent with chronic degenerative disease or interstitial nephrosis.
- Prominent, hypoechoic pancreas – The pancreatic changes are most consistent with mild pancreatitis or a recent episode of pancreatic inflammation.
- Mildly enlarged mesenteric lymph nodes – The prominent abdominal lymph nodes are most consistent with reactive lymphadenitis or lymphoid hyperplasia. Neoplastic infiltration is considered less likely.

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

There are no focal lesions to associate with the renal failure such as stones, masses, obstructions, etc. Being a Shar Pei, this pet should be screened for glomerulonephritis with a urine protein to creatinine ratio, provided urinalysis and culture are normal. Additionally, recommend blood pressure evaluation. If glomerulonephritis is present, then consider starting medical treatment and possibly treatment for amyloid(?), additionally looking for any additional sources of inflammation such as pyoderma, otitis, etc. that can exacerbate the kidney disease.

The left limb of the pancreas appears somewhat prominent. The significance of this is currently unclear. If there are any signs associated with pancreatitis, consider treatment. Additionally, you could consider evaluating the diet, and consider a lower fat option.



**PATIENT**

Scarlett Walls

**SPECIES**

Canine

**BREED**

Shar Pei

**SEX**

Spayed Female

**AGE**

8 Years

**WEIGHT**

44 Pounds

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

**IMAGING  
PERFORMED BY**

Sorbo

**HOSPITAL NAME**

Mill Brook AC

**REFERRING VET**


Dr. Jeffers

**INVOICE**

45769

**DATE**

3/8/23





**PATIENT**

Scarlett Walls

**SPECIES**

Canine

**BREED**

Shar Pei

**SEX**

Spayed Female

**AGE**

8 Years

**WEIGHT**

44 Pounds

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

**IMAGING  
PERFORMED BY**

Sorbo

**HOSPITAL NAME**

Mill Brook AC

**REFERRING VET**


Dr. Jeffers

**INVOICE**

45769

**DATE**

3/8/23



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

kathleen.sennello@sonopath.com