



**PATIENT**

Sushi Netto

**SPECIES**

Canine

**BREED**

Shih Tzu

**SEX**

Neutered Male

**AGE**

6 Months

**WEIGHT**

13.5 Pounds

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Shari Reffi, CVT

**HOSPITAL NAME**

Summit Dog & Cat  
Hospital

**REFERRING VET**

Dr. Lepkowski

**INVOICE**

43308

**DATE**

6/20/23

**PRESENTING CLINICAL SIGNS**

Frequent V+ started in May. Developed diarrhea early June. Fecal=neg. Current meds: Famotidine, Sucralfate, Metronidazole.

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**Urinary System**

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The prostate is normal in size (0.55 cm) and shape for this neutered male dog. The parenchyma is homogenous and the external margins are smooth. The prostatic urethra appears normal with no evidence of irregularity, invasion, mass effect or calculi.

The left kidney has a normal shape and size (3.54 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (3.89 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

**Adrenal Glands**

The left adrenal gland is normal in size measuring 0.30 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is normal in size measuring 0.44 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

**Spleen**

The spleen is subjectively normal in size, echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

**Liver**


The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is homogenous echotexture. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

The gallbladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and primarily anechoic. The cystic and common bile ducts are normal/not visible.

**Gastrointestinal**



<b>PATIENT</b>	The stomach contains minimal luminal contents. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.
Sushi Netto	
<b>SPECIES</b>	The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. Jejunum wall measures 0.25 cm. Duodenum wall measures 0.35 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.
Canine	
<b>BREED</b>	
Shih Tzu	The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.
<b>SEX</b>	
Neutered Male	<b>Pancreas</b>
<b>AGE</b>	The pancreas is normal and isoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.
6 Months	<b>Free Abdomen</b>
<b>WEIGHT</b>	Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.
13.5 Pounds	
<b>INTERPRETED BY</b>	<b>ULTRASONOGRAPHIC FINDINGS</b>
Beth Johnson, DVM DACVIM	<ul style="list-style-type: none"> <li>No significant ultrasonographic lesions visualized.</li> </ul>
<b>IMAGING PERFORMED BY</b>	<b>INTERPRETATION OF THE FINDINGS &amp; FURTHER RECOMMENDATIONS</b>
Shari Reffi, CVT	A focal lesion responsible for the reported vomiting and diarrhea is not clearly visualized. Unfortunately, there are many causes for vomiting that cannot be definitively diagnosed by ultrasound alone.
<b>HOSPITAL NAME</b>	Consider metabolic causes. Recommend full lab work, possibly including screening for Addison's disease +/- a liver function test (no evidence of a portosystemic shunt visualized, but it cannot 100% be ruled out).
Summit Dog & Cat Hospital	If metabolic disease is thought unlikely, then consider primary gastrointestinal causes of vomiting and diarrhea.
<b>REFERRING VET</b>	Consider such differentials as food allergy/dietary intolerance, GI parasitism, pancreatitis, dysbiosis, and much less likely IBD or GI neoplasia.
Dr. Lepkowski	<ul style="list-style-type: none"> <li>Consider a novel protein/hydrolyzed protein diet (exclusively at least 4-6 weeks)</li> <li>Consider a GI panel to Texas A&amp;M for evaluation of B12 levels, folate, PLI/TLI etc.. to further evaluate for pancreatic/small intestinal disease.</li> <li>Recommend screening for Addison's disease.</li> <li>Recommend chronic probiotic therapy.</li> <li>If not already done, recommend screening and empirical treatment for GI parasitism.</li> </ul>
<b>INVOICE</b>	
43308	
<b>DATE</b>	
6/20/23	If there is no response to these therapies and diagnostics and symptoms are persistent, consider reevaluation (lab work +/- repeat imaging) prior to considering endoscopic evaluation.
	Recommend three view thoracic radiographs to evaluate for possible concurrent thoracic disease/involvement.



**PATIENT**

Sushi Netto

**SPECIES**

Canine

**BREED**

Shih Tzu

**SEX**

Neutered Male

**AGE**

6 Months

**WEIGHT**

13.5 Pounds

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING PERFORMED BY**

Shari Reffi, CVT

**HOSPITAL NAME**

Summit Dog & Cat  
Hospital

**REFERRING VET**


Dr. Lepkowski

**INVOICE**

43308

**DATE**

6/20/23





**PATIENT**

Sushi Netto

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

**SPECIES**

Canine

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**BREED**

Shih Tzu

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

info@sonopath.com

**SEX**

Neutered Male

**AGE**

6 Months

**WEIGHT**

13.5 Pounds

**INTERPRETED BY**

Beth Johnson, DVM  
DACVIM

**IMAGING  
PERFORMED BY**

Shari Reffi, CVT

**HOSPITAL NAME**

Summit Dog & Cat  
Hospital

**REFERRING VET**

Dr. Lepkowski

**INVOICE**

43308

**DATE**

6/20/23