


**DATE PRESENTING CLINICAL SIGNS**

6/4/26

**Patient History:** CC: One episode of diarrhea and vomiting about 6-9 weeks ago. H: Healthy his entire life. PE: NSF

**PATIENT**

Joey McNemar

**Current Medications:** N/A.

**Labwork Results:** Labwork attached, reported as: Chem: WNL. CBC: WNL. T4: WNL. Idexx Spec FPL: 12 TAMU profile done about 3 weeks later, still showed an elevated spec cpl

**Date of Previous IntraPet Ultrasound:** No previous.

**SPECIES**

Feline

**Sedation:** Not required to complete full diagnostic ultrasound.

**Stat Report:** Not requested.

**Imaging Performed by:** Rachel Brillhart, RDMS.

**BREED**

DSH

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**
**Urinary System**

The urinary bladder is moderately distended with mild primarily suspended echogenic debris present. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or calculi. Echogenic debris of this type can be associated with small crystals, cellular debris and proteinaceous debris.

**AGE**

10/1/12

The left kidney has a normal shape and size (3.68 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

**WEIGHT**

13.6 lbs

The right kidney has a normal shape and size (4.02 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

**INTERPRETED BY**

Kathleen Sennello DVM,  
MS, Diplomate ACVIM  
(Small Animal Internal  
Medicine)

**HOSPITAL NAME**

Harborside Mobile  
Veterinary Clinic

**Adrenal Glands**

The left adrenal gland is normal in size measuring 0.42 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is normal in size measuring 0.43 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

**REFERRING VET**

Dr. Hawkins

**Spleen**

The spleen is subjectively normal in size (0.86 cm), echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

**INVOICE**

75708

**Liver**

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is homogenous echotexture. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

The gall bladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and likely incidental at this time. The cystic and common bile ducts are normal/not visible.

### ***Gastrointestinal***

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.36cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

The visualized areas of duodenum, jejunum and ileum have a uniform diameter with minimal fluid distension. Wall thickness is normal to slightly increased. Bowel loops follow a typical curvilinear path with distinct wall layering, but some areas display a prominent muscularis layer which does not display the typical 1:3 muscularis:mucosa layer ratio. Duodenum wall measures 0.32 cm. Jejunum wall measures 0.24 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

### ***Pancreas***

The pancreas is prominent and mottled in both limbs. There is no evidence of nodules or cystic lesions. Prominent pancreatic duct noted measuring 0.22 cm. There is very mild reactive mesentery noted in the region. No surrounding free fluid noted.

### ***Free Abdomen***

Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

## **ULTRASONOGRAPHIC FINDINGS**

- Mild suspended echogenic debris in the urinary bladder – The echogenic debris in the bladder lumen could be consistent with cells, crystals, and/or mucus.
- Pancreatic changes most consistent with chronic pancreatic remodeling/resolving pancreatitis. Chronic active pancreatitis is possible.
- Prominent muscularis layer of the small intestine – The small intestinal wall changes could be consistent with an underlying inflammatory process. These types of changes can sometimes be seen in normal older cats. Correlate with clinical signs.


## **INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**


The pancreas is prominent in both limbs with prominent pancreatic duct and some subtle surrounding inflammation on some views. Given the history provided, this could be consistent with resolving pancreatitis +/- mild smoldering pancreatitis. Often the pancreas will not return to normal appearance, so ongoing treatment is largely guided by clinical symptoms. In this situation I would consider a combination ultra low-fat/hydrolyzed protein prescription diet (Royal Canin has this) if the patient will tolerate it, or ideally a low-fat diet, and continued monitoring. Monitoring of PLI levels can be frustrating, as they do not always

correlate with clinical findings.

The small intestine appears diffusely “ropey”, possibly consistent with inflammatory type change. In the absence of chronic gastrointestinal symptoms and a normal GI panel, the significance of this is uncertain. Initially I would likely consider dietary management (hydrolyzed protein prescription diet) if possible or continued monitoring.

Pancreatic neoplasia is unlikely based on the presentation and appearance, although continued monitoring is warranted. If the patient is having relapses, etc., repeat imaging could be considered.





The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)  
info@sonopath.com