



PATIENT

Chico Wilson

SPECIES

Canine

BREED

Chihuahua

SEX

Neutered Male

AGE

4 Years

WEIGHT

8.2 kg

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING PERFORMED BY

Dr. Gira

HOSPITAL NAME

Resolution Vet
Ultrasound

REFERRING VET

Dr. Gira

INVOICE

38559

DATE

6/8/22

PRESENTING CLINICAL SIGNS

Chronic episodes of vomiting , inappetence and lethargy .
Abnormal PE/Chem/CBC/UA Results: Previous BW during the last episode (March 2022) Amylase 1,733 500 - 1,500 U/L Lipase 3,951 200 - 1,800 U/L Current episode occurred 2 days ago . Spec cPL run today before ultrasound was performed Spec cPL 73 0 - 200 µg/L

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The visualized areas of prostate and surrounding tissue appear normal. Unfortunately, the prostate is not fully visualized likely due to its intrapelvic location. Correlate with rectal exam findings.

The left kidney has a normal shape and size (4.42 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (4.65 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

Adrenal Glands

The left adrenal gland is normal in size measuring 0.58 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The right adrenal gland is normal in size measuring 0.52 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

Spleen

The spleen is subjectively normal in size, echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

Liver

The liver is large in size, and normal in echogenicity with smooth peripheral margins. The parenchyma is heterogenous in echotexture with subtle, indistinct focal mottling. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

The gall bladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. There is a moderate amount of non-organized echogenic debris. The cystic and common bile ducts are normal/not visible.



PATIENT

Gastrointestinal

Chico Wilson

The stomach is moderately dilated with fluid and irregular shadowing material most consistent with normal ingesta and gas. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layering is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

SPECIES

Canine

The visualized areas of duodenum, jejunum and ileum have a uniform diameter with minimal fluid distension. Wall appears subjectively, mildly increased. Bowel loops follow a typical curvilinear path with distinct wall layering. Duodenum wall measured 0.39 cm. Jejunum wall measured 0.31 cm.

BREED

Chihuahua

Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

SEX

Neutered Male

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

Pancreas

AGE

4 Years

The pancreas is prominent and mottled compared to the surrounding isoechoic mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

Free Abdomen

WEIGHT

8.2 kg

Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

ULTRASONOGRAPHIC FINDINGS

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

- Prominent, mottled pancreas – The pancreatic changes are most consistent with age-related parenchymal remodeling, potentially secondary to a prior inflammatory episode, early fibrosis or chronic pancreatitis.
- Large, heterogeneous liver – The diffuse hepatic changes are non-specific and could be consistent with vacuolar hepatopathy, nodular hyperplasia, inflammatory/immune-mediated disease, fibrosis, extramedullary hematopoiesis, toxic hepatopathy (e.g., copper), infiltrative neoplasia (less likely) or other hepatopathy.
- Moderate gallbladder debris – The significance of the aggregated gallbladder sludge is unclear. This could represent an early mucocele, cholestasis, or may be secondary to fasting.
- Subjectively thickened small intestine – The mild small intestinal wall changes may be a normal variant in this patient or could be consistent with an inflammatory process (e.g., inflammatory bowel disease).

IMAGING PERFORMED BY

Dr. Gira

HOSPITAL NAME

Resolution Vet
Ultrasound

REFERRING VET

Dr. Gira

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

No prominent focal lesions were visualized today associated with the gastrointestinal tract to explain the episodes of GI upset reported. Overall, the pancreas is mildly prominent, and for such a small dog, the GI tract appears slightly thickened, but there is no loss of layering, etc. Additionally, the liver appears somewhat heterogeneous. The significance of this is difficult to say with no liver enzyme elevation. Consider a liver function test to determine if the liver is of concern.

INVOICE

38559

DATE

6/8/22



PATIENT

Chico Wilson

- Consider an novel protein/hydrolyzed protein prescription diet. If the patient won't eat this, you can consult/hire a veterinary nutritionist to formulate a novel protein/hydrolyzed protein homemade diet, which they owner will prepare and may be more palatable.

SPECIES

Canine

- Consider chronic probiotic therapy.
- Consider a baseline cortisol level to rule out Addison's disease.

BREED

Chihuahua

- Consider pre- and post-prandial bile acids to rule out liver dysfunction.
- Recommend a GI panel to Texas A&M for a qualitative PLI, TLI, cobalamin and folate to further evaluate the pancreas and small intestine.

SEX

Neutered Male

- If GI signs persist despite dietary change and the above diagnostics, then consider obtaining GI biopsies.

AGE

4 Years

WEIGHT

8.2 kg



INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)




IMAGING PERFORMED BY

Dr. Gira

HOSPITAL NAME

Resolution Vet
Ultrasound



REFERRING VET

Dr. Gira

INVOICE

38559

DATE

6/8/22



PATIENT

Chico Wilson

SPECIES

Canine

BREED

Chihuahua

SEX

Neutered Male

AGE

4 Years

WEIGHT

8.2 kg

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Dr. Gira

HOSPITAL NAME

Resolution Vet
Ultrasound

REFERRING VET


Dr. Gira

INVOICE

38559

DATE

6/8/22



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)

kathleen.sennello@sonopath.com