



PATIENT

Sidney Angel

PRESENTING CLINICAL SIGNS

appetite decreased, elevated liver and pancreatic values. On meloxicam, gabapentin, famotidine.

SPECIES

Canine

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

BREED

Cairn Terrier

The left kidney has a normal shape and size (4.66 cm). Overall echogenicity is slightly hyperechoic with poor corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of perinephric inflammation or effusion. There is no evidence of pyelectasia, infarcts or hydroureter. There are numerous small non obstructive nephroliths/mineralizations present. Renal vasculature is normal.

SEX

Spayed Female

AGE

12 Years

The right kidney has a normal shape and size (4.67 cm). Overall echogenicity is slightly hyperechoic with poor corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of perinephric inflammation or effusion. There is no evidence of pyelectasia, infarcts or hydroureter. There are numerous small non obstructive nephroliths/mineralizations present. Renal vasculature is normal.

WEIGHT

18.6 Pounds

Adrenal Glands

The left adrenal gland is normal in size measuring 0.52 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

The right adrenal gland is normal in size measuring 0.61 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

IMAGING PERFORMED BY

Diane McFadden

Spleen

The spleen is subjectively normal in size and the echotexture is homogenous. The splenic capsule is smooth with no visible irregularities. Rare discrete focal hyperechoic, perivascular parenchymal abnormalities are present. The appearance of these lesions is most consistent with benign splenic myelolipomas. The blood flow through the hilus and splenic parenchyma appears normal.

HOSPITAL NAME

AH of Roxbury

Liver

The liver is subjectively normal/large in size. It is of mixed echogenicity with diffusely irregular peripheral margins. The parenchyma is severely heterogeneous and nodular. The visible portions of the vasculature and biliary tract appear normal. There are diffuse hypoechoic and hyperechoic foci throughout the liver, resulting in a diffusely nodular pattern.

REFERRING VET

Dr. Elia

INVOICE

24781

The gallbladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are primarily anechoic. The cystic and common bile ducts are normal/not visible.

DATE

8/18/21



PATIENT

Gastrointestinal

Sidney Angel

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

SPECIES

Canine

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. The duodenum measured as normal (between 0.3-0.5cm in wall thickness) and the jejunum measured as normal (between 0.2-0.47cm.)

BREED

Cairn Terrier

Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

SEX

Spayed Female

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

Pancreas

AGE

12 Years

The pancreas is prominent and hypoechoic as compared to the surrounding isoechoic mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

Free Abdomen

WEIGHT

18.6 Pounds

Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

ULTRASONOGRAPHIC FINDINGS

- Severely nodular liver – The diffuse hepatic changes are non-specific and could be consistent with vacuolar hepatopathy, nodular hyperplasia, inflammatory/immune-mediated disease, fibrosis, extramedullary hematopoiesis, toxic hepatopathy (e.g., copper), infiltrative neoplasia (less likely) or other hepatopathy.
- Prominent, hypoechoic pancreas – The pancreatic changes are most consistent with mild pancreatitis or a recent episode of pancreatic inflammation.
- Decreased corticomedullary distinction in both kidneys with non-obstructive nephroliths – Mild loss of corticomedullary distinction in both kidneys could be consistent with chronic degenerative disease or interstitial nephrosis.

IMAGING PERFORMED BY

Diane McFadden

HOSPITAL NAME

AH of Roxbury

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

REFERRING VET

Dr. Elia

The liver changes observed are dramatic, but not necessarily specific for disease process. These types of changes can be seen with diffuse neoplasia as well as severe regenerative nodules and nodular hyperplasia. I would lean towards this being a benign process, but the extent of damage to the liver is currently unknown. A fine needle aspirate is an excellent start. With some of these disease processes, a biopsy is necessary to obtain a definitive diagnosis. Recommend liver function testing, an fPLI with a GI panel to get more information on the pancreas and the GI tract, evaluation of coagulation parameters, and possibly supplemental therapy for the liver depending on what blood work changes are present. Although this could be a benign process, it could also be an indicator of end stage disease. Correlate with clinical findings.

INVOICE

24781

DATE

8/18/21



PATIENT

Sidney Angel

SPECIES

Canine

BREED

Cairn Terrier

SEX

Spayed Female

AGE

12 Years

WEIGHT

18.6 Pounds

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Diane McFadden

HOSPITAL NAME

AH of Roxbury

REFERRING VET





Dr. Elia

INVOICE

24781

DATE

8/18/21





PATIENT

Sidney Angel

SPECIES

Canine

BREED

Cairn Terrier

SEX

Spayed Female

AGE

12 Years

WEIGHT

18.6 Pounds

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Diane McFadden

HOSPITAL NAME

AH of Roxbury

REFERRING VET


Dr. Elia

INVOICE

24781

DATE

8/18/21



The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)
kathleen.sennello@sonopath.com