



PATIENT

Scarlet Montgomery

SPECIES

Canine

BREED

Boxer

SEX

Spayed Female

AGE

7 Years

WEIGHT

67 Pounds

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Dr. Jack Reese

HOSPITAL NAME

Willow Run VC

REFERRING VET

Dr. Jack Reese

INVOICE

25969

DATE

9/30/21

PRESENTING CLINICAL SIGNS

Chronic history of mild IBD (diagnosed via intestinal biopsy) responsive to dietary change. Recent history of increased water intake, possible anxiety at home - pacing, panting.
Abnormal PE/Chem/CBC/UA Results: Full lab work shows mild elevation in ALP (276) Urinalysis unremarkable LDDST pending

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

The left kidney has a normal shape and size (6.54 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

The right kidney has a normal shape and size (7.04 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

Adrenal Glands

The left adrenal gland is normal in size measuring 0.49 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

The region of the right adrenal (between right cranial kidney and vena cava) is unremarkable, but the adrenal is not distinctly visualized. No evidence of a mass effect.

Spleen

The spleen is subjectively normal in size, echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

Liver

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is mildly heterogenous in echotexture with subtle, indistinct focal mottling. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.


The gallbladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are primarily anechoic. The cystic and common bile ducts are normal/not visible.

Gastrointestinal

The stomach is moderately dilated with fluid and irregular shadowing material most consistent with normal ingesta and gas. It measures at a normal thickness of <0.7cm with some variability due to the



PATIENT	presence of rugal folds. The distinction of the gastric wall layering is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.
Scarlet Montgomery	
SPECIES	The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. Jejunum wall measured 0.27 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.
Canine	
BREED	The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.
Boxer	
SEX	Pancreas The pancreas is normal and isoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.
Spayed Female	
AGE	Free Abdomen Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.
7 Years	
WEIGHT	PRIMARY FINDINGS
67 Pounds	<ul style="list-style-type: none"> Mildly heterogeneous liver – The diffuse hepatic changes are non-specific and could be consistent with vacuolar hepatopathy, nodular hyperplasia, inflammatory/immune-mediated disease, fibrosis, extramedullary hematopoiesis, toxic hepatopathy (e.g., copper), infiltrative neoplasia (less likely) or other hepatopathy.
INTERPRETED BY	SECONDARY FINDINGS
Kathleen Sennello DVM, MS, Diplomate ACVIM (Small Animal Internal Medicine)	<ul style="list-style-type: none"> Mild ingesta in the gastric lumen – Correlate with feeding history. If patient was fasted, then consider delayed gastric emptying or partial obstruction (none observed).
IMAGING PERFORMED BY	INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS
Dr. Jack Reese	The ultrasonographic lesions observed were relatively subjective and mild. An obvious cause for the increased thirst and mild ALP elevation is not identified. Consider:
HOSPITAL NAME	<ul style="list-style-type: none"> Urinalysis and culture to rule out urinary tract infection causing PU/PD Confirm normal calcium levels Consider testing for Leptospirosis if this could be clinically relevant If clinical signs fit with Cushing's disease, you could consider adrenal function testing (low-dose Dexamethasone suppression test is pending per history) Consider quantitating water intake. If the patient is panting more at home, sometimes this will cause dry mouth and increase thirst. Consider environmental/behavioral factors. You could consider a liver function test to ensure that mild liver enzyme elevations do not represent impaired hepatic function. Consider 3-view thoracic radiographs to evaluate for any evidence of intrathoracic disease.
Willow Run VC	
REFERRING VET	
Dr. Jack Reese	
INVOICE	
25969	
DATE	
9/30/21	



PATIENT

Scarlet Montgomery

SPECIES

Canine

BREED

Boxer

SEX

Spayed Female

AGE

7 Years

WEIGHT

67 Pounds

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Dr. Jack Reese

HOSPITAL NAME

Willow Run VC

REFERRING VET


Dr. Jack Reese

INVOICE

25969

DATE

9/30/21



The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)
kathleen.sennello@sonopath.com