

PATIENT

Reba Regan

PRESENTING CLINICAL SIGNS

SPECIES

Canine

Sedated 0.05ml Dex/otrb IV- P presented on 7/6 for possible constipation, vomiting, panting/restless. P has been acting normally since discharge on the same day. P was treated with a bland diet and probiotics. LABs revealed elevated liver values- GI signs resolved and NSF findings on RADs taken that day. AUS for elevated ALT and ALP.

BREED

Australian Cattle Dog

Abnormal PE/Chem/CBC/UA Results: Summary of Laboratory Abnormalities: cbc - rbc 9.22 (5.5-8.5), hgb 19.9 (12-18), hct 62.77 (37-55), chem - alt 357 (10-118), alp 504 (20-150), glu 123 (60-110), cpl normal (not consistent with pancreatitis) RADS- NSF

SEX

Spayed Female

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

AGE

8 Years

The left kidney has a normal shape and size (5.89 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydronephrosis. Renal vasculature is normal.

WEIGHT

47.6 Pounds

The right kidney has a normal shape and size (6.1 cm). Overall echogenicity is normal with adequate corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of focal perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydronephrosis. Renal vasculature is normal.

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

Adrenal Glands

The left adrenal gland is normal in size measuring 0.59 cm at the caudal pole. It is observed in its normal position cranial to the left renal artery. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

IMAGING BY

Loetitia Saint-Jacques,
LVT

The right adrenal gland is normal in size measuring 0.54 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

HOSPITAL NAME

Desert Hills AH

Spleen

The spleen is subjectively normal in size, echotexture is homogenous, and the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

REFERRING VET

Dr. Michelle Caldwell

Liver

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is heterogenous in echotexture with subtle, indistinct focal mottling. The visible portions of the vasculature and biliary tract appear normal. There are too numerous to count, ill-defined, irregular, subtle hypoechoic nodules visualized, varying in size from approximately 0.5-1.5 cm. Examples measure 0.69, 0.78, 0.65, and 1.3 cm.

INVOICE

41088

DATE

9/6/22



PATIENT

Reba Regan

The gallbladder lumen is moderately distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. Luminal contents are mild and primarily anechoic. The cystic and common bile ducts are normal/not visible.

SPECIES

Canine

Gastrointestinal

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

BREED

Australian Cattle Dog

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. The duodenum measured as normal (between 0.3-0.5cm in wall thickness) and the jejunum measured as normal (between 0.2-0.47cm.) Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

SEX

Spayed Female

AGE

8 Years

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

WEIGHT

47.6 Pounds

Pancreas

The pancreas is prominent and mottled compared to the surrounding isoechoic mesentery. There is no evidence of nodules or cystic lesions. There is no evidence of regional mesenteric inflammation or fluid.

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

Free Abdomen

Evaluation of the peritoneal cavity did not reveal any evidence of effusion. The lymph nodes appear normal in size. The right sublumbar lymph node measures 0.79 cm. The left measures 0.80 cm. The omentum is of normal echogenicity.

Other

A brief view of the heart was submitted. No significant pericardial effusion was seen.

IMAGING BY

Loetitia Saint-Jacques,
LVT

ULTRASONOGRAPHIC FINDINGS

HOSPITAL NAME

Desert Hills AH

- Prominent, mottled pancreas – The pancreatic changes are most consistent with age-related parenchymal remodeling, potentially secondary to a prior inflammatory episode, early fibrosis or chronic pancreatitis.

REFERRING VET

Dr. Michelle Caldwell

- Heterogeneous liver with ill-defined hypoechoic nodules – The diffuse hepatic changes are non-specific and could be consistent with vacuolar hepatopathy, nodular hyperplasia, inflammatory/immune-mediated disease, fibrosis, extramedullary hematopoiesis, toxic hepatopathy (e.g., copper), infiltrative neoplasia (less likely) or other hepatopathy. The nodules observed trend toward a more benign process but underlying neoplasia cannot be ruled out.

INVOICE

41088

DATE

9/6/22



PATIENT

Reba Regan

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

SPECIES

No focal lesions are visualized today associated with the liver. The parenchyma is severely heterogeneous and there are numerous small, irregular, hypoechoic nodules visualized. Consider the following:

Canine

BREED

- Consider close evaluation of history for possible toxic changes examine medications, diet, dietary indiscretion etc...

Australian Cattle Dog

- Consider PCR on urine/serum for leptospirosis (if not on antibiotics)/serology if recent antibiotic history

SEX

- If not already done, consider pre and post prandial bile acids to evaluate liver function

Spayed Female

- Consider Fine needle aspirate if round cell neoplasia is on your differentia list (25 g needle, normal coags)

AGE

- If no response to supportive care (Denamarin, fluids, antibiotics, +/- ursodiol etc.) Consider liver biopsy with samples obtained for histopathology, culture, and copper levels.

8 Years

The pancreatic changes are extremely subtle and likely insignificant.

WEIGHT

47.6 Pounds

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING BY


Loetitia Saint-Jacques,
LVT

HOSPITAL NAME

Desert Hills AH

REFERRING VET

Dr. Michelle Caldwell




INVOICE

41088

DATE

9/6/22



PATIENT

Reba Regan

SPECIES

Canine

BREED

Australian Cattle Dog

SEX

Spayed Female

AGE

8 Years

WEIGHT

47.6 Pounds

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

IMAGING BY

Loetitia Saint-Jacques,
LVT

HOSPITAL NAME

Desert Hills AH

REFERRING VET


Dr. Michelle Caldwell

INVOICE

41088

DATE

9/6/22



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)
kathleen.sennello@sonopath.com