



PATIENT

Lily Addison

PRESENTING CLINICAL SIGNS

Patient has been lethargic, anorexic, having diarrhea, and vomiting for 5 days.
Abnormal PE/Chem/CBC/UA Results: Patient is icteric. ALP: 1611, ALT: unreadable, Tbili: 13.2

SPECIES

Canine

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

BREED

Golden Retriever X

The urinary bladder is moderately distended with anechoic urine. The Bladder wall, trigone, ureteral papillae and visible urethra (to a depth of 2cm) appear normal with no evidence of wall thickening, mucosal irregularities, masses or cystic calculi.

SEX

Spayed Female

The left kidney has a normal shape and size (5.3 cm). Overall echogenicity is slightly hyperechoic with poor corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

AGE

9 Years

The right kidney has a normal shape and size (6.0 cm). Overall echogenicity is slightly hyperechoic with poor corticomedullary distinction and a typical 1:3 cortex:medulla ratio. There is no evidence of perinephric inflammation or effusion. There is no evidence of pyelectasia, nephroliths, infarcts or hydroureter. Renal vasculature is normal.

WEIGHT

56.5

Adrenal Glands

The region of left adrenal (Cranial to left renal artery) is unremarkable but the adrenal is not distinctly visualized. No evidence of a mass effect.

The right adrenal gland is normal in size measuring 0.69 cm at the caudal pole. It is observed in its normal position between the cranial aspect of the right kidney and the caudal vena cava. It is normal in appearance (uniformly hypoechoic) and shape with no evidence of a mass effect.

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

Spleen

The spleen is subjectively normal in size. The spleen echotexture is heterogenous and mottled, the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized.

IMAGING PERFORMED BY

Dallas Ramberg

Liver

The liver is subjectively normal in size, and echogenicity with smooth peripheral margins. The parenchyma is heterogenous in echotexture with subtle, indistinct focal mottling. The visible portions of the vasculature and biliary tract appear normal. No focal nodules or cystic lesions are observed.

HOSPITAL NAME

Lone Mountain AH

REFERRING VET

Dr. Jacquelyn Shoppa

The gallbladder lumen is significantly distended. The wall of the gall bladder is not thickened and has a smooth mucosal surface. There is a large amount of non-organized, echogenic debris within the gallbladder. I am unable to clearly visualize the common bile duct.

Gastrointestinal

INVOICE

25291

The stomach contains minimal luminal contents. It measures at a normal thickness of <0.7cm with some variability due to the presence of rugal folds. The distinction of the gastric wall layers is adequate and there is no impression of reduced peristaltic activity. No masses or focal lesions were observed.

DATE

9/9/21

The visualized areas of duodenum, jejunum and ileum have a relatively uniform diameter with minimal fluid distension. Wall thickness is normal. Bowel loops follow a curvilinear path with distinct wall layering maintaining the typical 1:3 muscularis:mucosa layer ratio. Duodenum wall measured 0.4 cm.



PATIENT

Lily Addison

Jejunum wall measured 0.38 cm. Visualized peristalsis appears appropriate. There were no focal lesions consistent with obstruction or a mass effect observed.

SPECIES

Canine

The ileocecal junction was visualized and exhibited normal intact wall layering and is subjectively of normal thickness. Sections of colon are visualized with formed fecal material and gas shadowing distally. There is no observed focal or generalized colon wall thickening or loss of layering.

BREED

Golden Retriever X

Pancreas

The pancreas is large and hypoechoic to surrounding mesentery. There is no evidence of nodules or cystic lesions. There is evidence of regional mesenteric inflammation. Consistent with severe pancreatitis.

SEX

Spayed Female

Free Abdomen

Evaluation of the peritoneal cavity did not reveal any evidence of effusion, or subjective lymphadenomegaly. The Medial iliac nodes appear normal and there was no evidence of a caudal aortic thrombus at the bifurcation. The omentum is of normal uniform echogenicity.

AGE

9 Years

PRIMARY FINDINGS

WEIGHT

56.5

- Large, hypoechoic pancreas surrounded by hyperechoic mesentery – The pancreatic changes are most consistent with severe pancreatitis/pancreatic inflammation. Recommend fPLI testing and continued monitoring for improvement or possible development of a pancreatic abscess. Consider fine needle aspirate if not improving.
- Distended gallbladder – suspect secondary obstruction due to pancreatitis.
- Heterogeneous liver – The diffuse hepatic changes are non-specific and could be consistent with vacuolar hepatopathy, nodular hyperplasia, inflammatory/immune-mediated disease, fibrosis, extramedullary hematopoiesis, toxic hepatopathy (e.g., copper), infiltrative neoplasia (less likely) or other hepatopathy.

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

SECONDARY FINDINGS

IMAGING PERFORMED BY

Dallas Ramberg

- Decreased corticomedullary distinction in both kidneys – Mild loss of corticomedullary distinction in both kidneys could be consistent with chronic degenerative disease or interstitial nephrosis.
- The spleen is subjectively normal in size. The spleen echotexture is heterogenous and mottled, the splenic capsule is smooth with no irregularities. The blood flow through the hilus and splenic parenchyma appears normal. No focal parenchymal abnormalities are visualized. This subtle mottling favors a more benign differential.

HOSPITAL NAME

Lone Mountain AH

REFERRING VET

Dr. Jacquelyn Shoppa

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS


There is severe pancreatic inflammation present with no obvious obstruction seen or gallbladder disease seen, increasing suspicion for a post-hepatic obstruction of the bile duct with an inflamed pancreas. I cannot rule out a concurrent hepatopathy, so consider a fine needle aspirate of the liver as long as coagulation parameters are normal. Recommend aggressive therapy for pancreatitis with follow up ultrasounds to look evidence of abscess formation. In severe cases, a biliary stent may need to be placed to relieve the obstruction. Recommend 3-view thoracic radiographs.

INVOICE

25291

DATE

9/9/21



PATIENT

Lily Addison

SPECIES

Canine

BREED

Golden Retriever X

SEX

Spayed Female

AGE

9 Years

WEIGHT

56.5

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Dallas Ramberg

HOSPITAL NAME

Lone Mountain AH

REFERRING VET





Dr. Jacquelyn Shoppa


INVOICE

25291

DATE

9/9/21





PATIENT

Lily Addison

SPECIES

Canine

BREED

Golden Retriever X

SEX

Spayed Female

AGE

9 Years

WEIGHT

56.5

INTERPRETED BY

Kathleen Sennello DVM,
MS, Diplomate ACVIM
(Small Animal Internal
Medicine)

**IMAGING
PERFORMED BY**

Dallas Ramberg

HOSPITAL NAME

Lone Mountain AH

REFERRING VET


Dr. Jacquelyn Shoppa

INVOICE

25291

DATE

9/9/21



The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Kathleen Sennello DVM,MS, Diplomate ACVIM (Small animal Internal Medicine)
kathleen.sennello@sonopath.com