



## PATIENT

Mollie Ritter

## PRESENTING CLINICAL SIGNS

Newly acquired murmur; No symptoms  
Abnormal PE/Chem/CBC/UA Results: All BW WNL

## SPECIES

Canine

## ULTRASONOGRAPHIC EXAMINATION OF THE HEART & ABDOMEN

## BREED

Terrier Mix

CANINE	MR	TR	LA/AO	LA/AO	FS	EF	EPSS
<b>CARDIAC PARAMETERS</b>	<b>VMAX</b> (m/s)	<b>VMAX</b> (m/s)	(Boon method)	(Heart Base; Swe)	(%)	(%)	(cm)
<b>NORMAL PARAMETER</b>	4.5-5.5	<2.7	1.3	<1.6	28-40	40-100	<0.6
<b>PATIENT</b>			--	1.25	43.3	77.9	0.2
CANINE	HR	AV	PV	BODY WEIGHT	LA	LVIDd	LVIDs
<b>CARDIAC PARAMETERS</b>	(BPM)	<b>VMAX</b> (m/s)	<b>MAX</b> (m/s)	(kg)	2D short axis Base view (cm)	Avg; 2D and m-mode short axis (cm)	Avg; 2D and m-mode short axis (cm)
<b>NORMAL PARAMETER</b>	50-100	0.7-1.7	0.7-1.6	BELOW	BELOW	BELOW	BELOW
<b>PATIENT</b>	NM	NM	NM		3.0	3.0	

## SEX

SF

## AGE

8 Years

## WEIGHT

22

## INTERPRETED BY

R. McKenzie Daniel,  
DVM, DABVP  
(Canine and Feline)

## IMAGING PERFORMED BY

Tasha

## HOSPITAL NAME

Dillsburg VC

## REFERRING VET

Dr. Crow

## INVOICE

49366

## DATE

1-7-22

### Cardiac Presentation

The echocardiogram in this patient demonstrated normal **left atrial** size based on 3 separate methods of LA evaluation. The cranial and caudal **mitral** valve leaflets presented normal linear structure, extension in systole, and union in diastole with normal kinesis. The **left ventricle** presented thicknesses with linear contour and was not dilated nor restricted. The **myocardium** presented normal echogenicity without subjective evidence of significant fibrotic or ischemic disease. **Contractility** of the ventricular walls was adequate and in normal range for this patient evidenced by the fractional shortening measurement and subjective evaluation of the different regions of the myocardium. The **left ventricular outflow** tract demonstrated normal laminar flow and subjective structural integrity. The **right atrium** and auricle revealed normal size, structure and content. No evidence of masses was noted. **Tricuspid** valvular assessment demonstrated adequate linear morphology and kinesis. The **right ventricle** was of normal size (1/3 diameter of LV), chordae structure, myocardial echogenicity and thickness. **Pulmonary outflow** tract assessment revealed normal valve structure, laminar flow, and diameter (approx. 1:1 pa/ao ratio). No visible **pericardial** or free pleura fluid was noted. The cranial **mediastinum** and **pericardial and extra-cardiac regions** were free of masses in the visible window.



<b>PATIENT</b>	<b><i>Urinary System</i></b>
Mollie Ritter	The urinary bladder, trigone, cystourethral junction, and visible pelvic urethra to a depth of 3.0 cm exhibited normal thickness and tone. Anechoic urine was present in the lumen with no uroliths or sediment. The ureteral papillae were normal. The ureters were not visible which is normal. No evidence of inflammatory or neoplastic changes were noted.
<b>SPECIES</b>	
Canine	No evidence of pathology in the area of the aortic trifurcation.
<b>BREED</b>	
Terrier Mix	Normal size and margination was present in the kidneys. A normal 1:3 cortex / medulla ratio and normal corticomedullary definition were maintained. The echogenicity of the cortex was similar to or slightly less than normal liver parenchyma while the medulla echogenicity was hypoechoic to the cortex with no evidence of pelvic dilation. The left kidney measured 4.8 cm in length. The right kidney measured 4.6 cm in length.
<b>SEX</b>	<b><i>Adrenal Glands</i></b>
SF	The left adrenal gland was uniform in size and contour with a uniformly hypoechoic parenchyma. The left adrenal gland measured 0.59 cm width at the caudal pole.
<b>AGE</b>	
8 Years	The right adrenal gland was not definitively visualized.
<b>WEIGHT</b>	<b><i>Spleen</i></b>
22	The spleen exhibited a finely textured and homogenous parenchyma which was hyperechoic to the liver and renal cortical parenchyma. The capsule was smooth and regular without apparent expansion. The splenic vasculature at the hilus was normal in volume with no evidence of congestion or thrombosis. Acute to chronic inflammatory, neoplastic, or benign parenchyma changes were not noted.
<b>INTERPRETED BY</b>	<b><i>Liver / Gallbladder</i></b>
R. McKenzie Daniel, DVM, DABVP (Canine and Feline)	The liver was subjectively normal in size, structure, and contour. The liver parenchyma was uniform and hypoechoic to the spleen with a mild coarse echotexture. The hepatic and portal vasculature were normal in appearance without signs of congestion. The gallbladder was non-distended in size with thin walls and primarily anechoic luminal content. The cystic and common bile ducts were normal.
<b>IMAGING PERFORMED BY</b>	<b><i>Gastrointestinal</i></b>
Tasha	The stomach presented intact wall layering with a normal wall layer ratio. The lumen of the stomach was empty with no signs of ileus, obstruction or foreign material.
<b>HOSPITAL NAME</b>	
Dillsburg VC	The small intestine presented intact wall layering with 1:3 muscularis/mucosa ratio. The lumen of the small intestine was empty with no signs of ileus, obstruction or foreign material.
<b>REFERRING VET</b>	Normal visible colon wall layers were present with apparent formed feces in lumen.
Dr. Crow	<b><i>Pancreas</i></b>
<b>INVOICE</b>	The parenchyma of the left limb, body and right limb of the pancreas presented isoechoic to the adjacent omental fat. A normal curvilinear capsule contour of the pancreas was present. The visible pancreatic duct was normal. No signs of active inflammation or neoplastic disease was evident.
49366	<b><i>Free Abdomen</i></b>
<b>DATE</b>	No overt lymphadenopathy or peritoneal effusion was present.
1-7-22	



## PATIENT

Mollie Ritter

## ULTRASONOGRAPHIC FINDINGS

- Sonographically unremarkable abdomen.
- Overtly normal cardiac structure and function.

## SPECIES

Canine

## INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

No evidence of abdominal visceral pathology.

## BREED

Terrier Mix

Normal overall cardiac structure and function without definitive cause of the murmur identified. In the absence of volume changes (dehydration) or anemia, a physiologic or flow murmur, small flow abnormality, valvular insufficiency, i.e., mitral valve insufficiency, may be possible. Mitral valve insufficiency may be suspected if the murmur is primarily left sided. Regardless for the lack of left or right heart chamber enlargement as well as lack of other clinical issues such as systolic dysfunction indicate that the risk of complication is low. No indication for cardiac medications.

## SEX

SF

Conservative monitoring of the murmur is recommended. Recheck full echocardiogram may be considered in 6-12 months, sooner if clinical signs suggestive of heart disease initiate or if the murmur progresses.

## AGE

8 Years

## WEIGHT

22

## INTERPRETED BY

R. McKenzie Daniel,  
DVM, DABVP  
(Canine and Feline)

## IMAGING PERFORMED BY

Tasha

## HOSPITAL NAME

Dillsburg VC

## REFERRING VET


Dr. Crow


## INVOICE

49366

## DATE

1-7-22





**PATIENT**

Mollie Ritter

**SPECIES**

Canine

**BREED**

Terrier Mix

**SEX**

SF

**AGE**

8 Years

**WEIGHT**

22



**INTERPRETED BY**

R. McKenzie Daniel,  
DVM, DABVP  
(Canine and Feline)

**IMAGING PERFORMED BY**

Tasha

**HOSPITAL NAME**

Dillsburg VC

**REFERRING VET**

Dr. Crow

**INVOICE**

49366

**DATE**

1-7-22

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

R. McKenzie Daniel, DVM, DABVP (Canine / Feline Practice)  
info@SonoPath.com