

<b>PATIENT</b>	<b>PRESENTING CLINICAL SIGNS</b>
Linc Martinsen	continued weight loss, on hepatosupport supplementation. Had an abd US performed 5 months ago, revealed likely chronic hepatitis. Any worsening of chronic hepatitis or other abnormalities seen? Further recommendations for treatment and diagnostics. On Hepatosupport. Previous scan 5/18/21.
<b>SPECIES</b>	
Canine	ALP 479, ALT 356, Albumin 27, BUN 1.9, Cholesterol 4.8, Glucose 5.7
<b>BREED</b>	<b>ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN</b>
Doberman	<b>Urinary System</b>
<b>SEX</b>	The urinary bladder, trigone, and cystourethral junction exhibited normal thickness and tone. Mild to moderate, dependent mineral was present. The ureteral papillae were normal. The ureters were not visible which is normal. No evidence of inflammatory or neoplastic changes was noted.
MN	The area of the aortic trifurcation was free of pathology.
<b>AGE</b>	
6 years	Normal size and margination were present in the kidneys. A normal 1:3 cortex / medulla ratio and normal corticomedullary definition were maintained. The echogenicity of the cortex was similar to or slightly less than normal liver parenchyma while the medulla echogenicity was hypoechoic to the cortex with no evidence of pelvic dilation. The left kidney measured 8.4 cm in length.
<b>WEIGHT</b>	<b>Adrenal Glands</b>
30.8 kg	The left and right adrenal glands were not definitively visualized owing to patient size, conformation, and presence of free fluid.
<b>INTERPRETED BY</b>	<b>Spleen</b>
R. McKenzie Daniel, DVM, DABVP	The spleen exhibited a finely textured and homogenous parenchyma which was hyperechoic to the liver and renal cortical parenchyma. The capsule was smooth and regular without apparent expansion. The splenic vasculature at the hilus was normal in volume with no evidence of congestion or thrombosis. Acute to chronic inflammatory, neoplastic, or benign parenchyma changes were not noted.
<b>IMAGING PERFORMED BY</b>	<b>Liver/ Gallbladder</b>
Crystal Hill	The liver was normal in size with subtle asymmetrical hepatic capsule margins. Generalized nonuniform and variably echogenic parenchyma were present. No distinct hepatic masse or nodules were noted. The gallbladder was non-distended in size with thin walls and primarily anechoic luminal content. The cystic and common bile ducts were normal.
<b>HOSPITAL NAME</b>	<b>Gastrointestinal</b>
Queensway AH	The stomach presented intact wall layering with a normal wall layer ratio. The lumen of the stomach was empty with no signs of ileus, obstruction, or foreign material.
<b>REFERRING VET</b>	
Dr. Bilinsky	The small intestine presented intact wall layering with 1:3 muscularis/mucosa ratio. The lumen of the small intestine was empty with no signs of ileus, obstruction, or foreign material.
<b>INVOICE</b>	
12418	Normal visible colon wall layers were present with apparent formed feces in lumen.
<b>DATE</b>	
10/21/21	



**PATIENT** *Pancreas*

Linc Martinsen

The parenchyma of the left limb, body, and right limb of the pancreas presented isoechoic to the adjacent omental fat. A normal curvilinear capsule contour of the pancreas was present. The visible pancreatic duct was normal. No signs of active inflammation or neoplastic disease were evident.

**SPECIES**

Canine

**Free Abdomen**

Mild to moderate, primarily anechoic, subjectively acellular to mildly cellular peritoneal free fluid was present. The omentum was of uniform echogenicity. No overt lymphadenopathy was noted.

**BREED**

Doberman

Rapid view of the heart revealed no evidence of pericardial masses or effusion in the visible window.

**ULTRASONOGRAPHIC FINDINGS**

**SEX**

**Primary Findings**

MN

- Chronic hepatopathy - suspect probable progressive to possible emerging end-stage hepatopathy, given the sonographic presentation, chronic history of elevated hepatic enzymes, presence of peritoneal free fluid, decreased BUN level, and low normal albumin

**AGE**

6 years

- Urinary bladder mineral

- Mild to moderate peritoneal free fluid

**WEIGHT**

30.8 kg

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

Nonspecific chronic hepatitis (immune-mediated, infectious, or other), is suspected with potential for some degree of vacuolar hepatic changes, parenchymal remodeling, and / or early fibrosis / cirrhosis.

**INTERPRETED BY**

R. McKenzie Daniel,  
DVM, DABVP

Coagulation panel and bile acid testing may be considered. Hepatic sampling is required for a definitive diagnosis. Correlation with effusion analysis and cytology may be considered. Continued hepatosupportive medications with additional therapy based on the clinical impression of the patient would be appropriate. Guarded prognosis, given the likelihood of chronic hepatic failure.

**IMAGING PERFORMED BY**

Crystal Hill

**HOSPITAL NAME**

Queensway AH

**REFERRING VET**

Dr. Bilinsky

**INVOICE**

12418

**DATE**

10/21/21





**PATIENT**

Linc Martinsen

**SPECIES**

Canine

**BREED**

Doberman

**SEX**

MN

**AGE**

6 years

**WEIGHT**

30.8 kg

**INTERPRETED BY**

R. McKenzie Daniel,  
DVM, DABVP

**IMAGING PERFORMED BY**

Crystal Hill

**HOSPITAL NAME**

Queensway AH

**REFERRING VET**


Dr. Bilinsky

**INVOICE**

12418

**DATE**


10/21/21



The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

R. McKenzie Daniel, DVM, DABVP (Canine / Feline Practice)  
info@SonoPath.com



## PATIENT

Linc Martinsen

## Canine Liver Disease & Treatment Recommendations

## SPECIES

Canine

<http://www.sonopath.com/K9LiverDisease>

## BREED

Doberman

**Description:** The etiologic causes of canine hepatic disease are vast and varied. Some cases may progress fairly rapidly, while others will remain static for a considerable length of time or even eventually reverse. Regardless of the cause, management is crucial to maintaining and optimizing quality of life. If possible, practitioners should obtain and be guided by a pathologic diagnosis so they can administer a treatment attuned to the underlying disease and arrive at a more exact prognosis.

## SEX

MN

**Dietary Management:** A lower protein diet to support liver dysfunction should be initiated, especially in cases where hepatic encephalopathy is also present. Since dietary protein is low, the protein quality and bioavailability must conversely be high. It should be noted that a protein-restricted diet is not appropriate in all cases of hepatic disease, especially during the early phases, as protein restriction is unnecessary when there are no signs of significant hepatic dysfunction.

## AGE

6 years

## WEIGHT

30.8 kg

Therapeutic diets, such as Hill's® i/d® and Royal Canin® Hepatic™, are excellent choices and contain enhanced levels of nutrients such as, but not limited to:

- Branched chain amino acids, which bypass liver metabolism and are used directly for skeletal muscle accretion.
- Vitamin E, which helps minimize and reduce oxidative damage and stress from free radicals produced by stressed hepatocytes.
- Vitamin B complex, which helps drive intermediary metabolism.
- Reduced copper.
- Extremely digestible protein sources with high biologic values, which help minimize the total amount of dietary protein needed and thus reduce blood ammonia levels.
- Carnitine, which helps drive fatty acids into the mitochondria for beta-oxidation and positive cellular energy balance.

## INTERPRETED BY

R. McKenzie Daniel,  
DVM, DABVP

## IMAGING PERFORMED BY

Crystal Hill

## HOSPITAL NAME

Queensway AH

## REFERRING VET

Dr. Bilinsky

**Medical Management:** The following list of medications is commonly used in the management of various hepatopathies or in the face of hepatic failure; however, each patient should be managed as an individual, and not all of the medications listed here are appropriate for each animal. One must always consider the definitive diagnosis of one's patient when developing a therapeutic plan. What follows is an outline of medical management recommendations for cholangiohepatitis and inflammatory hepatopathy/chronic hepatitis.

## INVOICE

12418


### Cholangiohepatitis

#### 1. Broad-spectrum antibiotics


## DATE

10/21/21


a) Amoxicillin: Give 20 mg/kg BID or amoxicillin/clavulanic acid (13.75 mg/kg PO BID) for potential suppurative hepatitis. Options: ampicillin: 20 mg/kg IV TID; cephalexin: 20 mg/kg IV or PO TID; enrofloxacin: 2.5-5 mg/kg PO BID if cholangiohepatitis is present or to decrease




<b>PATIENT</b>	ammonia production; gentamycin: 2 mg/kg TID IM or SC for 5-7 days if sepsis or peritonitis is present. Monitor renal function if aminoglycosides are utilized.
Linc Martinsen	
<b>SPECIES</b>	b) Metronidazole: Give 10-20 mg/kg BID in combination with amoxicillin/clavulanic acid or enrofloxacin for cholangiohepatitis because of its efficacy against anaerobic bacteria and/or for its immunomodulating effects. The dose is decreased to 7.5 mg/kg PO TID in the face of hepatic failure and/or encephalopathy. Controls ammonia production in the colon, decreases bacteria absorbed through portal circulation, and reduces cell-mediated immune responses (anti-inflammatory properties).
Canine	
<b>BREED</b>	
Doberman	2. <i>Hepatic support</i>
<b>SEX</b>	a) S-adenosylmethionine (SAME): Give 20 mg/kg/day PO on an empty stomach (1-2 hours before feeding). It is available in 90 mg tablets that are not to be broken. SAME replenishes glutathione and aids in cellular detoxification; it also has anti-arthritis effects. SAME is an anti-inflammatory and antioxidant. It also promotes hepatocellular regeneration and rectifies RBC membrane abnormalities in dogs with liver disease or oxidative damage.
MN	
<b>AGE</b>	b) Milk Thistle: Administer as silybin or silymarin extracts (a high-quality supplement is essential). Acts as an antioxidant and free radical scavenger; decreases hepatotoxin binding; improves glutathione concentrations; aids in iron chelating; and promotes choleresis. Give 5-15 mg/kg/day PO.
6 years	
<b>WEIGHT</b>	c) Ursodiol (Actigall): Give 10-15 mg/kg PO once daily, with food, to stimulate bile flow and decrease cholestasis. Tablets (250 mg) or capsules (300 mg) are available; however, ursodiol can also be compounded into a liquid to dose small patients. It has immunomodulatory, anti-fibrotic, and choleric effects, anti-copper storage benefits, and stabilizes mitochondrial function.
30.8 kg	
<b>INTERPRETED BY</b>	d) Vitamin E: Must be coupled with good nutrition and other antioxidants to avoid accumulation of tocopheroxyl radicals. To that end, supplementation with SAME may help ensure that adequate GSH (mitochondrial glutathione) concentrations are achieved. Give 10-15 IU/kg/day PO (100-400 IU) in a water-soluble form twice daily, as well as with Vitamin C 25 mg/kg/day.
R. McKenzie Daniel, DVM, DABVP	
<b>IMAGING PERFORMED BY</b>	e) Cobalamin and Thiamine (B12 and B1): Give 250ug SC weekly.
Crystal Hill	
<b>HOSPITAL NAME</b>	<b>Inflammatory hepatopathy/chronic hepatitis</b>
Queensway AH	1. <i>Immunosuppressive agents</i>
<b>REFERRING VET</b>	a) Prednisone or prednisolone: Administer if inflammatory disease has been diagnosed by biopsy, beginning at 2 mg/kg/day for 2-4 weeks; subsequently reduce to 1 mg/kg/day. Once remission has been achieved, taper to 0.5 mg/kg/day (or to the lowest tolerable dose) over 2-4 weeks. Steroids may be discontinued if a different immunosuppressive medication is effective at controlling inflammation (i.e., azathioprine or cyclosporine) since they are contraindicated with hepatic encephalopathy. Possible negative sequelae of corticosteroids include increased water retention and potentiation of gastrointestinal ulceration. In the face of portal hypertension and ascites, dexamethasone is preferred—it does not exhibit mineralocorticoid activity and thus does
Dr. Bilinsky	
<b>INVOICE</b>	
12418	
<b>DATE</b>	
10/21/21	



<b>PATIENT</b>	not potentiate water retention as compared to prednisone—at 0.2-0.4 mg/kg orally once daily. Taper in a similar manner.
Linc Martinsen	
<b>SPECIES</b>	b) Azathioprine (Imuran): Give 50 mg/m <sup>2</sup> /day or 2 mg/kg/day as a long-term alternative to prednisone. The dose can be decreased to 1 mg/kg and eventually given every other day if there is a positive response. Check CBC and platelet count biweekly for the first 2 months and then monthly thereafter. Taper every 2-4 weeks to the lowest effective dose while monitoring transaminase levels. It can often be dosed on alternate days to prednisone. Possible negative side effects include bone marrow suppression and hepatic necrosis. Cyclosporine has been proposed as an alternative immunosuppressant in the management of chronic hepatitis and may allow one to cease concurrent steroid therapy; however, this has not been thoroughly investigated as of yet.
Canine	
<b>BREED</b>	
Doberman	
<b>SEX</b>	2. <i>Hepatic Support</i>
MN	See medications listed in the previous section.
<b>AGE</b>	3. <i>Anti-fibrotics</i>
6 years	a) Colchicine: Give 0.03 mg/kg/day. Colchicine acts as an anti-inflammatory agent, stabilizes membranes, and stimulates collagenase production, thereby diminishing fibrosis. Colchicine should be used to treat hepatic fibrosis based on biopsy results; however, it can also be considered when ascites is present, and when hepatic fibrosis and cirrhosis are highly suspected based on sonographic appearance and clinical findings. It can result in adverse effects, including vomiting, diarrhea, and inappetance. Discontinue until clinical signs resolve, and reinstitute at a lower dose and up-titrate slowly.
<b>WEIGHT</b>	
30.8 kg	
<b>INTERPRETED BY</b>	4. <i>Hepatic Encephalopathy</i>
R. McKenzie Daniel, DVM, DABVP	a) Lactulose: Give 0.5 ml/kg orally 2-3 times daily to soften the stool. It helps manage hepatic encephalopathy by combining with ammonium in the GI tract and thus decreasing circulating ammonia levels. Use in conjunction with low dose metronidazole. Lactulose can also be given as a retention enema in an encephalopathic crisis.
<b>IMAGING PERFORMED BY</b>	b) Metronidazole: Give at 7.5 mg/kg PO TID. Neomycin is an alternative and can be administered at 22 mg/kg PO BID-TID.
Crystal Hill	
<b>HOSPITAL NAME</b>	c) L-Carnitine: Give 200-400 mg/day. Normally synthesized by the liver, L-Carnitine enhances ammonia elimination and is indicated in cases of hepatic encephalopathy and lipidosis. Carnitine must be in the L-form.
Queensway AH	
<b>REFERRING VET</b>	5. <i>Copper Chelation</i>
Dr. Bilinsky	Use chelation when copper toxicity has been documented on biopsy and quantification has been performed to confirm toxic levels.
<b>INVOICE</b>	
12418	
<b>DATE</b>	a) D-penicillamine: Give 10-15 mg/kg PO BID on an empty stomach. This is a copper chelator and should only be used based on a quantitative analysis of copper. Possible side effects include vomiting and inappetance. Do not give in conjunction with zinc.
10/21/21	



<b>PATIENT</b>	b) 2,3,2 Tetramine (Syprine, Cuprid): Give 5-7 mg/kg PO BID on an empty stomach (1-2 hours before eating). An alternative to D-penicillamine for those dogs that are intolerant.
Linc Martinsen	
<b>SPECIES</b>	c) Zinc gluconate, acetate, or sulfate (acetate is best tolerated): Give 15-10 mg/kg elemental zinc divided BID for 2-6 months as a loading dose. Administer on an empty stomach (30-60 minutes before eating). Reduce to half the dose during the maintenance phase. A low copper diet is preferred (i.e., therapeutic diets, such as Hill's I/d® or Royal Canin® Hepatic™, are advisable). Zinc binds with intestinal copper to avoid absorption in the gastrointestinal tract and may be used alone in mild cases of copper toxicity. The goal is to reach zinc serum levels of 200-600 ug/dl; levels should initially be measured every 2-3 months. Give this medication on an empty stomach or with tuna fish to avoid vomiting. Zinc is not as effective as D-penicillamine and is only used in mild cases. It is not used in conjunction with D-penicillamine.
Canine	
<b>BREED</b>	
Doberman	
<b>SEX</b>	6. Portal Hypertension and Ascites
MN	
<b>AGE</b>	a) Spironolactone: If ascites is present secondary to portal hypertension, spironolactone can be dosed at 1-2 mg/kg PO BID; it is the diuretic of choice. Alternatively, spironolactone can be used in conjunction with furosemide (0.5-1 mg/kg PO BID) or hydrochlorothiazide; one should administer 1 mg/kg PO BID if given in conjunction with another diuretic. Monitor renal function and electrolytes diligently.
6 years	
<b>WEIGHT</b>	b) Famotidine: Give 0.5 mg/kg PO BID in cases of portal hypertension that result in gastrointestinal bleeding/melena.
30.8 kg	
<b>INTERPRETED BY</b>	<b>General Notes on Therapeutic Management:</b> Given that a primary function of the liver is to metabolize oral medications via the portal system (first past effect), numerous medications may result in higher systemic exposure to parent compounds in the face of hepatic insufficiency or failure. Drugs that are inactivated by the liver, produce hepatic damage, or require hepatic metabolism should be avoided. These include: lincomycin, clindamycin, streptomycin, chloramphenicol, sulfonamides, erythromycin, hetacillin, phenobarbital, diazepam, oxy- or chloro-tetracyclines, azole antifungals, nonsteroidal anti-inflammatory drugs (NSAIDs), theophylline or chloramphenicol, combinations of cimetidine and metronidazole, and combinations of enrofloxacin and theophylline or cisapride. In cases of hepatic lipidosis, glucocorticoids, anabolic steroids, and lipotropic agents containing methionine should be avoided as they result in the production of encephalopathic toxins (metacarpans). Glucocorticoids are indicated for cholangitis, but only after lymphoma and hepatic lipidosis have been ruled out.
R. McKenzie Daniel, DVM, DABVP	
<b>IMAGING PERFORMED BY</b>	
Crystal Hill	
<b>HOSPITAL NAME</b>	
Queensway AH	
<b>REFERRING VET</b>	<b>References:</b>
Dr. Bilinsky	Bauer E. Hepatic disease, nutritional therapy, and the metabolic environment. <i>J Am Vet Med Assoc</i> 1996;209(11):1850-54.
<b>INVOICE</b>	Bradley AM and Twedt DC. Cyclosporine therapy for canine chronic hepatitis: a retrospective study. Proceedings from the American College of Veterinary Internal Medicine, Anaheim, CA, June 15-18, 2011.
12418	
<b>DATE</b>	Center SA. Chronic hepatitis, cirrhosis, breed-specific hepatopathies, copper storage hepatopathy, suppurative hepatitis, granulomatous hepatitis, and idiopathic hepatic fibrosis. In:
10/21/21	



**PATIENT**

Guilford WG, Center SA, Strombeck DR, Williams DA, and Meyer DJ, eds: *Strombeck's Small Animal Gastroenterology 3rd ed.* Philadelphia, PA: WB Saunders; 1996:705-65.

Linc Martinsen

**SPECIES**

Center S. Metabolic, antioxidant, nutraceutical, probiotic, and herbal therapies relating to the management of hepatobiliary disorders. *Vet Clin North Am Small Anim Pract* 2004;34(1):67-172.

Canine

**BREED**

Center SA. Diagnostic and managerial pearls for selected hepatobiliary disorders. Proceedings from the American College of Veterinary Internal Medicine. New Orleans, LA, May 30-June 2, 2012.

Doberman

**SEX**

Davenport D. Antimicrobial therapy for gastrointestinal, pancreatic, and hepatic disorders. *Probl Vet Med* 1990;2(2):374-93.

MN

**AGE**

Hackett T, Twedt D, Gustafson D. Milk thistle and its derivative compounds: a review of opportunities for treatment of liver disease. *J Vet Intern Med* 2013;27(1):10-16.

6 years

Thompson M, Meyer D, Senior D. Effects of treatment with ursodeoxycholic acid on bile acid profiles in a dog with chronic hepatic disease. *J Vet Intern Med* 1991;5(2):130.

**WEIGHT**

30.8 kg

**INTERPRETED BY**

R. McKenzie Daniel,  
DVM, DABVP

**IMAGING  
PERFORMED BY**

Crystal Hill

**HOSPITAL NAME**

Queensway AH

**REFERRING VET**

Dr. Bilinsky

**INVOICE**

12418

**DATE**

10/21/21