

PATIENT

Brownie Santos

SPECIES

Canine

BREED

Cockapoo

SEX

Male Neutered

AGE

10y

WEIGHT

33.2 lbs

INTERPRETED BY

R. McKenzie Daniel,
 DVM, DABVP
 (Canine and Feline)

IMAGING PERFORMED BY

Becca Hamilton

HOSPITAL NAME

Englewood VC

REFERRING VET

Dr. Ezik

INVOICE

13166

DATE

2/6/26

PRESENTING CLINICAL SIGNS

History:

- Follow up 6 month on invoice 10882
- Benign hepatopathy
- nonorganized gallbladder debris (non-mucocele)
- Splenomegaly with non-disruptive cranial splenic nodule
- static mild chronic renal changes
- Meds: Denamarin, Ursodiol

Abnormal PE/Chem/CBC/UA Results: ALP 314

ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN

Urinary System

The urinary bladder, trigone, cystourethral junction, and visible pelvic urethra to a depth of 3.0 cm exhibited normal thickness and tone. Anechoic urine was present in the lumen with no uroliths or sediment, mineral or calculi. The ureteral papillae were normal. The ureters were not visible which is normal. No evidence of inflammatory or neoplastic changes were noted.

The area of the residual prostate appeared normal and free of pathology.

The area of the aortic trifurcation was free of pathology.

Normal size and margination was present in the kidneys. A normal 1:3 cortex / medulla ratio was maintained. The medulla and cortices were uniform in texture with some increased echogenicity and mild loss of corticomedullary symmetry and definition expected for the age of the patient. No evidence of pelvic dilation was present. The left kidney measured 5.7 cm in length. The right kidney measured 6.1 cm in length.

Adrenal Glands

The left adrenal gland was uniform in size and contour with a uniformly hypoechoic parenchyma. The left adrenal gland measured 0.52 cm width at the caudal pole. The right adrenal gland was uniform in size and contour with a uniformly hypoechoic parenchyma. The right adrenal gland measured 0.52 cm width at the caudal pole.

Spleen

Previously noted non-capsule deforming splenic nodule measuring 0.83 cm in diameter. The spleen was normal in size and appearance compared to the previous study without evidence of progression.

Liver

The liver presented mildly enlarged in size. The parenchyma of the liver was subjectively normal in echogenicity compared to the spleen and renal cortices. The liver parenchyma was uniform with a mildly coarse echotexture. The capsule of the liver was symmetrically rounded to mildly swollen in margination. The hepatic and portal vasculature were normal in appearance without signs of congestion. The gallbladder was non distended in size with mild, non-organized, echogenic,



PATIENT

Brownie Santos

SPECIES

Canine

BREED

Cockapoo

SEX

Male Neutered

AGE

10y

WEIGHT

33.2 lbs

INTERPRETED BY

R. McKenzie Daniel,
DVM, DABVP
(Canine and Feline)

IMAGING PERFORMED BY

Becca Hamilton

HOSPITAL NAME

Englewood VC

REFERRING VET

Dr. Ezik

INVOICE

13166

DATE

2/6/26

nonmineralized biliary sludge. The cystic duct and common bile ducts were normal without evidence of dilation.

Gastrointestinal

The stomach presented intact wall layering with a normal wall layer ratio. The lumen of the stomach was empty with no signs of ileus, obstruction or foreign material.

The small intestine presented intact wall layering with 1:3 muscularis/mucosa ratio. The lumen of the small intestine was empty with no signs of ileus, obstruction or foreign material.

Normal visible colon wall layers were present with apparent formed feces in lumen.

Pancreas

The parenchyma of the left limb, body and right limb of the pancreas presented isoechoic to the adjacent omental fat. A normal curvilinear capsule contour of the pancreas was present. The visible pancreatic duct was normal. No signs of active inflammation or neoplastic disease was evident.

Free Abdomen


No overt lymphadenopathy or peritoneal effusion was present.

ULTRASONOGRAPHIC FINDINGS

- Static benign hepatopathy
- Static mild, non-organized gallbladder debris (non-mucocele)
- Static splenic nodule
- Static mild chronic renal changes

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

No evidence of progressive previously noted pathology. Continued sonographic monitoring and current hepato-supportive medications are recommended.





PATIENT

Brownie Santos

SPECIES

Canine

BREED

Cockapoo

SEX

Male Neutered

AGE

10y

WEIGHT

33.2 lbs

INTERPRETED BY

R. McKenzie Daniel,
 DVM, DABVP
 (Canine and Feline)

IMAGING PERFORMED BY

Becca Hamilton

HOSPITAL NAME

Englewood VC

REFERRING VET


Dr. Ezik

INVOICE

13166

DATE

2/6/26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

R. McKenzie Daniel, DVM, DABVP (Canine / Feline Practice)

info@sonopath.com



PATIENT

Brownie Santos

SPECIES

Canine

BREED

Cockapoo

SEX

Male Neutered

AGE

10y

WEIGHT

33.2 lbs

INTERPRETED BY

R. McKenzie Daniel,
DVM, DABVP
(Canine and Feline)

**IMAGING
PERFORMED BY**

Becca Hamilton

HOSPITAL NAME

Englewood VC

REFERRING VET

Dr. Ezik

INVOICE

13166

DATE

2/6/26