



PATIENT

Erin Verlezza

SPECIES

Feline

BREED

DSH

SEX

FS

AGE

11

WEIGHT

9.5

INTERPRETED BY

R. McKenzie Daniel,
DVM, DABVP
(Canine and Feline)

IMAGING PERFORMED BY

Jenn

HOSPITAL NAME

Rockaway AH

REFERRING VET

Dr. Salazar

INVOICE

10689

DATE

3/18/26

PRESENTING CLINICAL SIGNS

History:

- Hx of hyperthyroidism bloody stool Flow murmur suspected

Abnormal PE/Chem/CBC/UA Results: ALT 247 ProbBnp normal

ULTRASONOGRAPHIC EXAMINATION OF THE HEART & ABDOMEN

FELINE CARDIAC PARAMETERS	BODY WEIGHT	HR (BPM)	IVSd (cm)	LVIDd (cm)	LVWd (cm)	FS (%)	EF (%)
NORMAL PARAMETER	-----	150-240	0.3-0.6	1.0-2.1	0.25-0.6	35-67	80-100
PATIENT	9.5	178	0.48	1.57	0.49	55	86
FELINE CARDIAC PARAMETERS	LA/AO (M-mode)	LA/AO HEART BASE (Sisson)	LAD LA MAX 4 Chamber		LVOT VEL. (m/s)	RVOT VEL. (m/s)	IVRT (m/)
NORMAL PARAMETER	<1.5	1.6	0.7-1.7		<1.6	<1.3	40-60
PATIENT	1.1	1.1	1.2		-	0.95	-
Adapted from June Boon, Veterinary Echocardiography, 1998 Sisson D et al. JVIM 1991; 5: 232, Jacobs et al. Am J Vet Res 1985; 46:1705							

Cardiac Presentation

The echocardiogram in this patient demonstrated normal **left atrial** size based on 3 separate LA measurements. The cranial and caudal **mitral** valve leaflets presented normal linear structure and kinetics. The **left ventricle** presented normal thicknesses with linear contour and was not dilated nor restricted. The **myocardium** presented normal echogenicity without subjective evidence of significant fibrotic or ischemic disease. **Contractility** of the ventricular walls was adequate and in normal range for this patient evidenced by the fractional shortening measurement and subjective evaluation of the different regions and angles of the myocardium. The **left ventricular outflow** tract demonstrated normal laminar flow and subjective structural integrity. The **right atrium** and auricle revealed normal size, structure and content. No evidence of masses was noted or chamber overload. **Tricuspid** valvular assessment demonstrated adequate linear morphology and kinetics. The **right ventricle** was of normal size (1/3 diameter of LV), chordae structure, myocardial echogenicity and thickness. **Pulmonic** tract assessment revealed normal valve structure, laminar flow, and diameter (approx. 1:1 pa/ao ratio). No visible **pericardial** or free pleura fluid was noted or extra cardiac pathology in the visible planes. The cranial **mediastinum and pericardial regions** were free of masses in the visible window.

Urinary System

The urinary bladder, trigone, cystourethral junction, and visible pelvic urethra to a depth of 3.0 cm exhibited normal thickness and tone. Anechoic urine was present in the lumen with no evidence of urine



PATIENT	or lumen sediment, mineral, or calculi. The ureteral papillae were normal. The ureters were not visible which is normal. No evidence of inflammatory or neoplastic changes was noted.
Erin Verlezza	No evidence of pathology in the area of the aortic trifurcation.
SPECIES	Normal size and margination were present in the kidneys. A normal 1:3 cortex / medulla ratio was maintained. The medulla and cortices were uniform in texture with some increased echogenicity and mild loss of corticomedullary symmetry and definition expected for the age of the patient. No evidence of pelvic dilation was present. The left kidney measured 3.9 cm in length. The right kidney measured 3.9 cm in length.
Feline	
BREED	
DSH	<i>Adrenal Glands</i>
SEX	No overt pathology was noted in the area of the left or right adrenal glands.
FS	<i>Spleen</i>
AGE	The spleen exhibited a finely textured and homogenous parenchyma which was hyperechoic to the liver and renal cortical parenchyma. The capsule was smooth and regular without apparent expansion. The splenic vasculature at the hilus was normal in volume with no evidence of congestion or thrombosis. Acute to chronic inflammatory, neoplastic, or benign parenchyma changes were not noted.
11	
WEIGHT	<i>Liver/ Gallbladder</i>
9.5	The liver was subjectively normal in size, structure, and contour. The liver parenchyma was uniform and hypoechoic to the spleen with a mild coarse echotexture. The hepatic and portal vasculature were normal in appearance without signs of congestion. The gallbladder was non-distended in size with thin walls and primarily anechoic luminal content. The cystic and common bile ducts were normal.
INTERPRETED BY	<i>Gastrointestinal</i>
R. McKenzie Daniel, DVM, DABVP (Canine and Feline)	The stomach presented intact wall layering with a normal wall layer ratio. The lumen of the stomach was empty without evidence of retained ingesta, fluid, or foreign material.
IMAGING PERFORMED BY	The small intestine presented intact wall layering with 1:3 muscularis/mucosa ratio. The lumen of the small intestine was empty with no signs of ileus, obstruction, or foreign material.
Jenn	Normal visible colon wall layers were present with apparent formed fecal matter in lumen.
HOSPITAL NAME	<i>Pancreas</i>
Rockaway AH	The pancreas was normal in size and contour with isoechoic to heterogeneous parenchyma compared to adjacent omentum. No signs of active inflammation or neoplasia. Suspect, distal left limb, thinly walled pancreatic cyst containing anechoic fluid was present, measuring 0.81 cm in diameter.
REFERRING VET	<i>Free Abdomen</i>
Dr. Salazar	No significant or swollen mesenteric lymphadenopathy was visualized. No evidence of peritoneal effusion was present.
INVOICE	
10689	
DATE	
3/18/26	



PATIENT

Erin Verlezza

SPECIES

Feline

BREED

DSH

SEX

FS

AGE

11

WEIGHT

9.5

ULTRASONOGRAPHIC FINDINGS

- Normal cardiac structure / function
- Mild heterogeneous pancreas with subjective distal left pancreatic cyst
- Mild age-related renal changes
- Sonographically normal gastrointestinal tract / colon with current formed fecal matter

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

There is no evidence of structural or functional cardiomyopathy. If a murmur is present, a benign flow murmur is probable. A small non-visualized flow abnormality cannot be definitively excluded. Regardless of classification, and if present, the hemodynamic effects of any murmur are low. There is no indication for cardiac medications. Recheck echocardiogram is suggested in 6-12 months, sooner if potential increase in murmur intensity or if clinically indicated.

A spec fPL could be considered to assess for chronic pancreatitis. Given reported blood stool, gastrointestinal support and empirical therapy for potential low-grade colitis is recommended.

INTERPRETED BY

R. McKenzie Daniel,
DVM, DABVP
(Canine and Feline)

IMAGING PERFORMED BY

Jenn

HOSPITAL NAME

Rockaway AH

REFERRING VET


Dr. Salazar

INVOICE

10689

DATE

3/18/26





PATIENT

Erin Verlezza

SPECIES

Feline

BREED

DSH

SEX

FS

AGE

11

WEIGHT

9.5

INTERPRETED BY

R. McKenzie Daniel,
DVM, DABVP
(Canine and Feline)

IMAGING PERFORMED BY

Jenn

HOSPITAL NAME

Rockaway AH

REFERRING VET


Dr. Salazar

INVOICE

10689

DATE

3/18/26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

R. McKenzie Daniel, DVM, DABVP (Canine/Feline Practice)
info@sonopath.com