



<b>PATIENT</b>	<b>PRESENTING CLINICAL SIGNS</b>
Bronx Kodack	diarrhea, not eating/drinking lethargy
<b>SPECIES</b>	<b>ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN</b>
Canine	<b>Urinary System</b>
<b>BREED</b>	The urinary bladder, trigone, cystourethral junction, and visible pelvic urethra exhibited normal thickness and tone. Anechoic urine was present in the lumen with no uroliths or sediment. The ureteral papillae were normal. The ureters were not visible which is normal. No evidence of inflammatory or neoplastic changes were noted.
Frenchie	Normal size and margination were present in the kidneys. A normal 1:3 cortex / medulla ratio and normal corticomedullary definition were maintained. The echogenicity of the cortex was similar to or slightly less than normal liver parenchyma while the medulla echogenicity was hypoechoic to the cortex with no evidence of pelvic dilation. The left kidney measured 4.7 cm in length. The right kidney measured 5.1 cm in length.
<b>SEX</b>	<b>Adrenal Glands</b>
MI	The left adrenal gland was uniform in size and contour with a uniformly hypoechoic parenchyma. The left adrenal gland measured 0.45 cm width at the caudal pole and 0.53 cm width at the cranial pole. The right adrenal gland was uniform in size and contour with a uniformly hypoechoic parenchyma. The right adrenal gland measured 0.62 cm width at the caudal pole.
<b>AGE</b>	<b>Spleen</b>
6mo	The spleen exhibited a finely textured and homogenous parenchyma which was hyperechoic to the liver and renal cortical parenchyma. The capsule was smooth and regular without apparent expansion. The splenic vasculature at the hilus was normal in volume with no evidence of congestion or thrombosis. Acute to chronic inflammatory, neoplastic, or benign parenchyma changes were not noted.
<b>WEIGHT</b>	<b>Liver/Gallbladder</b>
25	The liver was subjectively normal in size, structure, and contour. The liver parenchyma was uniform and hypoechoic to the spleen with a mild coarse echotexture. The hepatic and portal vasculature were normal in appearance without signs of congestion. No evidence of portosystemic vascular anomaly. The gallbladder was non-distended in size with thin walls and primarily anechoic luminal content. The cystic and common bile ducts were normal.
<b>INTERPRETED BY</b>	<b>Gastrointestinal</b>
R. McKenzie Daniel, DVM, DABVP (Canine and Feline)	The stomach presented intact wall layering with a normal wall layer ratio. The lumen of the stomach was empty with no signs of ileus, obstruction or foreign material.
<b>IMAGING PERFORMED BY</b>	The small intestine presented intact wall layering with 1:3 muscularis/mucosa ratio. The lumen of the small intestine was empty potential mild increased gas pattern with no signs of ileus, obstruction or foreign material.
Jenn	Normal visible colon wall layers were present with apparent formed to shadowing feces in lumen.
<b>HOSPITAL NAME</b>	<b>Pancreas</b>
Rockaway Animal Hospital	
<b>REFERRING VET</b>	
Dr. Maniar	
<b>INVOICE</b>	
13248ag	
<b>DATE</b>	
03/21/2023	



**PATIENT**

Bronx Kodack

The parenchyma of the left limb, body and right limb of the pancreas presented isoechoic to the adjacent omental fat. A normal curvilinear capsule contour of the pancreas was present. The visible pancreatic duct was normal. No signs of active inflammation or neoplastic disease was evident.

**SPECIES**

Canine

**Free Abdomen**

No omental masses, overt lymphadenopathy or peritoneal effusion was present.

**ULTRASONOGRAPHIC FINDINGS**

**BREED**

Frenchie

- Sonographically unremarkable abdomen.

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

**SEX**

MI

Overall, no overt evidence of significant abdominal visceral specifically GI pathology as a definitive cause of the patient's clinical signs. No evidence of gastroenterocolic obstructive pattern or foreign material. Structurally insignificant inflammatory bowel episode, enterotoxic insult, occult parasitism or other inflammatory gastroenteropathy are all potentials. Hospitalization with as needed supportive care which may include IVF, gastroprotectants +/- antibiotic as well as empirical deworming should prove beneficial. Although considered unlikely considering normal adrenal presentation, a resting cortisol level to rule out occult Addison's disease could be considered given clinical signs and lack of abdominal visceral pathology.

**AGE**

6mo

**WEIGHT**

25

**INTERPRETED BY**

R. McKenzie Daniel,  
DVM, DABVP  
(Canine and Feline)

**IMAGING PERFORMED BY**


Jenn

**HOSPITAL NAME**

Rockaway Animal  
Hospital

**REFERRING VET**

Dr. Maniar



**INVOICE**

13248ag

**DATE**

03/21/2023



**PATIENT**

Bronx Kodack

**SPECIES**

Canine

**BREED**

Frenchie

**SEX**

MI

**AGE**

6mo

**WEIGHT**

25

**INTERPRETED BY**

R. McKenzie Daniel,  
DVM, DABVP  
(Canine and Feline)

**IMAGING PERFORMED BY**

Jenn

**HOSPITAL NAME**

Rockaway Animal  
Hospital

**REFERRING VET**


Dr. Maniar

**INVOICE**

13248ag

**DATE**

03/21/2023



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

R. McKenzie Daniel, DVM, DABVP (Canine/Feline Practice)  
[mac.daniel@sonopath.com](mailto:mac.daniel@sonopath.com)