



PATIENT	PRESENTING CLINICAL SIGNS
Dewi McNeill-Bindon	has recently been seen for limping - ran labwork to monitor carprofen, proteinuria present
SPECIES	Current Medications carprofen - d/c until after imaging
Canine	Radiographic Findings none Primary Question/Differential to Be Answered in This Exam r/o kidney diz
BREED	Abnormal PE/Chem/CBC/UA Results: proteinuria 3+ UPC 3.1
Corgi	ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN
SEX	<i>Urinary System</i>
MN	The urinary bladder, trigone, cystourethral junction, and visible pelvic urethra to a depth of 4 cm exhibited normal thickness and tone. Anechoic urine was present in the lumen with no uroliths or sediment. The ureteral papillae were normal. The ureters were not visible which is normal. No evidence of inflammatory or neoplastic changes were noted.
AGE	
11yr	Normal size and margination were present in the kidneys. A normal 1:3 cortex / medulla ratio was maintained. The medulla and cortices were uniform in texture with some increased echogenicity and mild loss of corticomedullary symmetry and definition expected for the age of the patient. No evidence of pelvic dilation was present. Bilateral pinpoint medullary mineral was present. The left kidney measured 6.1 cm in length. The right kidney measured 6.8 cm in length.
WEIGHT	
38lb	The area of the iliac trifurcation was free of pathology including no evidence of medial, iliac or sublumbar lymphadenopathy or masses. No evidence of distal aortic thrombus.
INTERPRETED BY	
R. McKenzie Daniel, DVM, DABVP (Canine and Feline)	The area of the prostate appeared normal and free of pathology; the prostate measured 0.81 cm in diameter.
IMAGING PERFORMED BY	<i>Adrenal Glands</i>
Jenna Walsh CVT	The left adrenal gland was uniform in size and contour with a uniformly hypoechoic parenchyma. The left adrenal gland measured 0.66 cm width at the caudal pole and 2.5 cm length. The right adrenal gland was uniform in size and contour with a uniformly hypoechoic parenchyma. The right adrenal gland measured 0.65 cm width at the caudal pole and 2.0 cm length.
HOSPITAL NAME	<i>Spleen</i>
VCA Vitality Animal Hospital	The spleen exhibited a finely textured and homogenous parenchyma which was hyperechoic to the liver and renal cortical parenchyma. The capsule was smooth and regular without apparent expansion. The splenic vasculature at the hilus was normal in volume with no evidence of congestion or thrombosis. Acute to chronic inflammatory, neoplastic, or benign parenchyma changes were not noted.
REFERRING VET	
Dr. Burgt	<i>Liver/Gallbladder</i>
INVOICE	
13685ag	The liver was subjectively normal in size, structure, and contour. The liver parenchyma was uniform and hypoechoic to the spleen with a mild coarse echotexture. The hepatic and portal vasculature were normal in appearance without signs of congestion. The gallbladder was non-distended in size with thin walls and primarily anechoic luminal content with minor non-organized echogenic debris. The cystic and common bile ducts were normal.
DATE	
05/01/2023	



PATIENT *Gastrointestinal*

Dewi McNeill-Bindon

The stomach presented intact wall layering with a normal wall layer ratio. The lumen of the stomach was empty with no signs of ileus, obstruction or foreign material.

SPECIES

Canine

The small intestine presented intact wall layering with 1:3 muscularis/mucosa ratio. The lumen of the small intestine was empty with no signs of ileus, obstruction or foreign material.

BREED

Corgi

Normal visible colon wall layers were present with apparent formed feces in lumen.

SEX

MN

Pancreas

The parenchyma of the left limb, body and right limb of the pancreas presented isoechoic to the adjacent omental fat. A normal curvilinear capsule contour of the pancreas was present. The visible pancreatic duct was normal. No signs of active inflammation or neoplastic disease was evident.

AGE

11yr

Free Abdomen

No omental masses, overt lymphadenopathy or peritoneal effusion was present.

WEIGHT

38lb

ULTRASONOGRAPHIC FINDINGS

- Non-specific mild chronic renal changes.
- Sonographically unremarkable urinary bladder.
- Normal liver.
- Gallbladder debris (non-mucocele).

INTERPRETED BY

R. McKenzie Daniel,
DVM, DABVP
(Canine and Feline)

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The increased UPC level is consistent with non-specific glomerulopathy with considerations including glomerulonephritis, amyloidosis or other. Infectious disease serology is suggested if clinically indicated. A renal surgical biopsy is required for a definitive diagnosis.


IMAGING PERFORMED BY

Jenna Walsh CVT

Empirical therapy for protein losing nephropathy which may include ACE inhibitor therapy/angiotensin receptor blocker and protein restricted diet as well as antiplatelet drugs is warranted. Monitoring of UPC is suggested.

HOSPITAL NAME

VCA Vitality Animal
Hospital



REFERRING VET

Dr. Burgt

INVOICE

13685ag

DATE

05/01/2023



PATIENT

Dewi McNeill-Bindon

SPECIES

Canine

BREED

Corgi

SEX

MN

AGE

11yr

WEIGHT

38lb

INTERPRETED BY

R. McKenzie Daniel,
DVM, DABVP
(Canine and Feline)

IMAGING PERFORMED BY

Jenna Walsh CVT

HOSPITAL NAME

VCA Vitality Animal
Hospital

REFERRING VET


Dr. Burgt

INVOICE


13685ag

DATE

05/01/2023



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not



PATIENT visible in the image/video clips provided.

Dewi McNeill-Bindon

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

SPECIES

R. McKenzie Daniel, DVM, DABVP (Canine / Feline Practice)

Canine

info@SonoPath.com

BREED

Corgi

Protein-Losing Nephropathy (PLN)

SEX

<http://www.sonopath.com/PLN>

MN

AGE

11yr

Description: Protein-losing nephropathy (PLN) is a common form of renal disease that typically affects dogs in middle age; it occurs less commonly in cats. Glomerular causes of renal protein loss encompass two broad categories: glomerulonephritis (GN) and amyloidosis. (The causes of GN in human medicine are more specifically differentiated based on a combination of histopathology, immunofluorescence, and electron microscopy findings.) Membranoproliferative glomerulonephritis is the most common cause of GN in dogs and is associated with infectious disease with secondary immune complex deposition as well as Lyme disease. Membranous nephropathy is the second most common cause of GN in dogs and the most common cause in cats. It occurs due to primary immune complex deposition on the urinary side of the basement membrane of the glomerulus, resulting in the leakage of albumin. Amyloidosis is caused by the deposition of amyloid A proteins in a β -pleated sheet configuration in the glomeruli. It is a familial disease in the Shar Pei but occurs as a reactive disease in other canine breeds. It is also inheritable in the Abyssinian cat, but the amyloidosis occurs in the medulla and is therefore not a protein-losing condition in this breed.

WEIGHT

38lb

INTERPRETED BY

R. McKenzie Daniel,
DVM, DABVP
(Canine and Feline)

Glomerular lesions can be associated with:

IMAGING PERFORMED BY

Jenna Walsh CVT

- Infectious diseases:
 - o Protozoan: Babesia, Hepatozoon, and Leishmania.
 - o Bacterial: Borrelia, Bartonella, Brucella, Ehrlichia, Mycoplasma, pyometra, pyoderma, endocarditis, and pyelonephritis.
 - o Viral: FeLV, FIV, and FIP.
 - o Fungal
 - o Helminthic: Dirofilaria.
- Non-infectious inflammatory diseases: pancreatitis, chronic dermatitis, inflammatory bowel disease, periodontal disease, polyarthritis, and systemic lupus erythematosus (SLE).
- Neoplasia: lymphoma, leukemia, and mast cell disease.
- Familial conditions in the soft-coated Wheaten Terrier, Shar Pei, Beagle, Cocker Spaniel, and Bernese Mountain dog.
- Idiopathic conditions.

HOSPITAL NAME

VCA Vitality Animal
Hospital

REFERRING VET

Dr. Burgt

INVOICE


13685ag

DATE

05/01/2023

Post-glomerular causes, such as hemorrhage and inflammation, also contribute to urine protein quantification.

Proteinuria Classifications: Patients can be divided into three tiers, depending on their clinical




PATIENT	characteristics:
Dewi McNeill-Bindon	Tier 1A: persistent subclinical proteinuria Tier 1B: persistent proteinuria with hypertension
SPECIES	
Canine	Tier 2A: proteinuria and hypoalbuminemia Tier 2B: proteinuria, hypoalbuminemia, and hypertension
BREED	
Corgi	Tier 3A: proteinuria and azotemia Tier 3B: proteinuria, azotemia, and hypertension Tier 3C: proteinuria, azotemia, hypertension, and hypoalbuminemia
SEX	
MN	
AGE	
11yr	
WEIGHT	
38lb	
INTERPRETED BY	
R. McKenzie Daniel, DVM, DABVP (Canine and Feline)	
IMAGING PERFORMED BY	
Jenna Walsh CVT	
HOSPITAL NAME	
VCA Vitality Animal Hospital	
REFERRING VET	
Dr. Burgt	
INVOICE	
13685ag	
DATE	
05/01/2023	

Diagnostics: Traditionally, urine protein loss has been detected either through a qualitative test, such as a urine dipstick, or with a semi-quantitative test, such as a urine protein-creatinine (UPC) ratio. When the latter is greater than 0.5, it is considered abnormal. False positive results can occur due to contamination of urine with red blood cells, white blood cells, and bacterial protein. Thus, one must use a urine sample with inactive sediment and a negative culture for measurement purposes. A 24-hour urine protein quantification is more accurate but technically more difficult to obtain, as it requires hospitalization and 24-hour urinary catheterization with a closed collection system. Pooling urine samples can be considered in cases where urine protein loss is stable. One must obtain three different urine samples, combine 1 ml from each sample to submit for a UPC test, and ensure that inactive sediments are present in all the samples. There should be a high degree of correlation between the UPC on the pooled sample and the mean of the three samples measured independently. Research has not yet demonstrated the accuracy of pooled samples for urine samples with high protein loss (i.e., in cases where the UPC is > 8).


Further diagnostic tests will depend on the tier classification. Once proteinuria is documented repeatedly, additional tests can be considered to assess for potential underlying causes, and, further to that, possible sources of antigen stimulation. Depending on presentation, tests may include:

- CBC and biochemical profile
- Urine culture and sensitivity
- 4DX
- Blood pressure measurement
- Thoracic and abdominal radiographs
- Spinal radiographs to assess for discospondylitis
- Abdominal ultrasound to assess for evidence of underlying infection or neoplasia
- Echocardiogram to assess for vegetative endocarditis and possible effects of hypertension
- Screen for Cushing's disease, especially if hypertensive (LDDST or ACTH stimulation)
- ANA
- Expanded tick or infectious disease screen
- Renal biopsy to differentiate among specific causes of PLN

Renal biopsy should be considered if proteinuria is severe (UPC > 3.5) and hypoalbuminemia and/or hypertension have been documented. Renal biopsy is an invasive procedure and should be considered only to determine if there is an underlying disease process that would benefit from specific therapy. If the patient is debilitated, severely azotemic, or has uncontrolled hypertension or coagulation



PATIENT	abnormalities, then the risk of the procedure and anesthesia may be too great and should not be pursued.
Dewi McNeill-Bindon	
SPECIES	Tissue samples should be submitted for a combination of light microscopy (in formalin; use with special stains), immunofluorescence (in Michel's solution or frozen), and electron microscopy (in formalin with glutaraldehyde). It is imperative to request special media before obtaining the biopsy. Samples can be obtained via ultrasound guidance, laparotomy, or laparoscopy, but cortical samples must be divided so that they can be placed in the three different media. One must ensure that the pre-surgical clotting profile and platelet count are both normal. Patients should undergo pre-biopsy and post-biopsy diuresis.
Canine	
BREED	
Corgi	
SEX	Treatment: The main goals of therapy are to i) reduce proteinuria (i.e., UPC < 1.0); ii) prevent a thrombotic event; iii) manage hypertension; and iv) replace fluid deficits. Fluid therapy should be approached cautiously, especially in patients with nephrotic syndrome. Standard therapy for PLN includes a low-protein diet, which in itself will reduce proteinuria, and the administration of an angiotensin-converting enzyme (ACE) inhibitor, such as enalapril (0.5 mg/kg PO BID) or benazepril (0.5 mg/kg PO Q24hr). Newer proposed therapeutic protocols include increasing the ACE inhibitor dose slowly while monitoring BUN and creatinine carefully. The dose can be raised to 1 mg/kg PO BID if needed, provided creatinine has not increased more than 30% from the baseline level.
MN	
AGE	
11yr	
WEIGHT	Another class of drugs currently being used is angiotensin receptor blockers, such as Losartan (the dose in azotemic dogs is 0.125-0.25 mg/kg/day PO Q12-24hr and 0.5-1.0 mg/kg/day in non-azotemic patients). This can be combined with an ACE inhibitor, but it is important to monitor BUN, creatinine, and potassium levels. Spironolactone has been used in people in combination with the other two classes of drugs to further modify the renin-angiotensin-aldosterone system (RAAS) (1-2 mg/kg PO BID); however, the effect of using all three drug classes in dogs has not yet been fully investigated. All of these medications are potassium sparing; thus, monitoring for hyperkalemia is important.
38lb	
INTERPRETED BY	Hypertension is managed with amlodipine (0.1-0.2 mg/kg PO Q12-24hr) when an ACE inhibitor is insufficient to control blood pressure. Supplementing with an anti-thrombotic agent, such as aspirin (1 mg/kg PO Q24hr), may be considered in advanced cases, especially once the patient is hypoalbuminemic. Omega-3 fatty acids can be given (0.25-0.5 g/day), but are typically increased in standard kidney diets.
R. McKenzie Daniel, DVM, DABVP (Canine and Feline)	
IMAGING PERFORMED BY	The most recent controversy in the management of glomerular diseases is the use of immunosuppressive medications. Because it is possible to arrive at a more definitive diagnosis in human patients, the use of immunosuppressive agents can be useful in the management of the disease, specifically when the disease is immune-mediated in its pathogenesis, such as SLE, membranous nephropathy, and minimal change disease glomerulonephritis. The procurement of a renal biopsy is being advocated in dogs so that practitioners can identify the population of patients that may benefit most from immunosuppressive therapy. Presently, there is no evidence-based medicine to suggest that immunosuppressive therapy should definitely be incorporated into a daily protocol for canine patients; however, it could be beneficial in some cases and may even result in remission. Further investigation is warranted. Trials are currently being conducted in patients with Lyme nephritis that are treated with immunosuppressive agents in addition to standard antibiotic therapy. The IRIS Treatment of Canine Glomerular Disease Study Group has suggested the trial use of immunosuppressive therapy in severe, persistent, or progressive PLN, even without a biopsy diagnosis in specific cases that are unresponsive
Jenna Walsh CVT	
HOSPITAL NAME	
VCA Vitality Animal Hospital	
REFERRING VET	
Dr. Burgt	
INVOICE	
13685ag	
DATE	
05/01/2023	



PATIENT	to standard therapy (i.e., nephrotic syndrome, progressively azotemic, hypoalbuminemic patients). One can also consider administering the following drugs: pulse steroid therapy, mycophenolate, cyclosporinamide, azathioprine, and chlorambucil. One should monitor blood work, UPC ratio, and blood pressure weekly for 2 weeks, then biweekly for 6 weeks, then monthly. If there is further deterioration, immunosuppressive therapy should be discontinued.
Dewi McNeill-Bindon	
SPECIES	References:
Canine	
BREED	Goldstein R and Polzin D. Treatment of canine glomerular disease: report of the IRIS treatment of canine glomerular disease study group. Proceedings from the American College of Veterinary Internal Medicine, Denver, CO, June 15-18, 2011.
Corgi	
SEX	Grauer GF, Greco DS, Getzy DM, et al. Effects of enalapril versus placebo as a treatment for canine idiopathic glomerulonephritis. J Vet Intern Med 2000;14:526-33.
MN	
AGE	Less GE, Cianciolo RE, and Clubb FJ. Renal biopsy and pathologic evaluation of glomerular disease. Top Companion Anim Med 2011;26(3):143-53.
11yr	
WEIGHT	LeVine DA, Zhang D, Vaden SL. The use of pooled vs. serial urine samples to measure urine protein:creatinine ratios. Vet Clin Pathol 2010;39(1):53-56.
38lb	
INTERPRETED BY	Nabity MB, Boggess MM, Kashtan CE, et al. Day-to-day variability in the urine protein:creatinine ratio in female dogs with stable glomerular proteinuria caused by x-linked hereditary nephropathy. J Vet Intern Med 2007;21:425-30.
R. McKenzie Daniel, DVM, DABVP (Canine and Feline)	Vaden SL. Glomerular Disease. Top Companion Anim Med 2011;26(3):128-34.
IMAGING PERFORMED BY	Vaden SL. Glomerular Diseases. In: Ettinger SJ and Feldman EC, eds. Textbook of Veterinary Internal Medicine, 7th Ed. St Louis, MI: Saunders Elsevier; 2010;2021-36.
Jenna Walsh CVT	Vaden SL. Microalbuminuria: What is it and how do I interpret it. Proceedings from the American College of Veterinary Internal Medicine, Charlotte, NC, June 4-7, 2003.
HOSPITAL NAME	Vaden SL et al. Urinary tract inflammation has a variable effect in urine albumin concentration. J Vet Intern Med 2002;16:378 (abstract).
VCA Vitality Animal Hospital	
REFERRING VET	
Dr. Burgt	
INVOICE	
13685ag	
DATE	
05/01/2023	