



PATIENT PRESENTING CLINICAL SIGNS

Rocco Figuera Reason for Visit: adr History: ~7y old M French Bulldog presents ADR. O reports yesterday pet did not eat, drink, and have a BM, pet would yelp when touched on R hip. Walking gingerly. O has had Rocco since August, someone gave him away without prev records.

SPECIES Abnormal PE/Chem/CBC/UA Results: CV/Respiratory: Normal heart rate and rhythm, no murmur, pulses strong and synchronous, normal bronchovesicular sounds. EENT: Clear AU. OU: lenticular sclerosis. No nasal discharge. No cough on tracheal palpation. Oral cavity: Mild to moderate dental tartar Musculoskeletal: BCS = 5/9. Ambulatory x 4, Stiff on hindlimbs - hyperextension of stifles. Normal ROM on exam. Uro/Perineum: Intact male, 2 descended testicles Abd/GI: Soft, non-painful. No masses or fluid wave palpated Lymph Nodes: No peripheral lymphadenopathy Neurological: Alert and appropriate. Possible neck pain, stiff on flexion to the sides. No CP deficits. Skin: Good hair coat. No ectoparasites seen Mentation: BAR Hydration: N 1.) CBC: WBC 17.26 (5.05-16.76), NEU 13.69 (2.95-11.64), EOS 0.01 (0.06-1.23), PLT 656 (148-484) 2) CHEM: ALT 146 (10-125)

BREED

French Bulldog

SEX

Male

RADIOGRAPHIC STUDY OF THE PELVIS & STIFLES

Lateral view of the pelvis with lateral projection of the stifles and ventrodorsal view of the pelvis totaling 2 images available for review.

AGE

7 Years

RADIOGRAPHIC FINDINGS

The patellae appear to be in situ in both stifle joints. No significant articular swelling is seen.

INTERPRETED BY

Nele Eley, DVM
Dr. med. Vet. DipECVDI

Cartilage remnants appear to be present in the tibial tuberosities which is an incidental finding.

The muscle volume of both hind limbs is symmetric and adequate.

The coxofemoral joints present within normal limits. No evidence of dysplasia or osteoarthritis is seen. There is no evidence of traumatic osseous injury or aggressive bone lesions.

HOSPITAL NAME

DPC Veterinary
Hospital

The coccygeal spine is represented by a series of hemi and block vertebrae and significantly shortened which is typical in corkscrew tailed breeds.

RADIOGRAPHIC DIAGNOSIS

REFERRING VET

Dr. Rivera

- Radiographically normal presentation of the stifles and coxofemoral joints.

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

No evidence of articular pathology of the stifle or coxofemoral joints is seen in either of the hind limbs. Early articular injury and patella luxation cannot be ruled out and require correlation with the clinical palpation.

INVOICE

49492

DATE

1-11-22



PATIENT

Rocco Figuera

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

SPECIES

Canine

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

BREED

French Bulldog

Nele Eley, DVM, Dr. med. vet., DipECVDI
European Specialist in Veterinary Diagnostic Imaging, Cert. Radiology,
Senior lecturer University of Giessen, Germany, Veterinary Faculty, Department of Radiology
Nele.Eley@sonopath.com

SEX

Male

AGE

7 Years

INTERPRETED BY

Nele Eley, DVM
Dr. med. Vet. DipECVDI

HOSPITAL NAME

DPC Veterinary
Hospital

REFERRING VET

Dr. Rivera

INVOICE

49492

DATE

1-11-22