



**PATIENT**

Ruby Morgan

**PRESENTING CLINICAL SIGNS**

Hx of right patellar sx 2 years ago. After pet was left with pet sitter for about a week pet became lame from right rear, O did find out that pet was jumping in and out of sitters car. P.E. unremarkable other than dental Dz and obesity, no pain elicited on manipulation of limb or no LROM coxofemoral joints. Pet did improve after a single PO dose of Meloxicam

**SPECIES**

Canine

**RADIOGRAPHIC STUDY OF THE STIFLES & PELVIS**

Mediolateral views of both stifles and ventrodorsal hip extended view of the pelvis totaling 3 images available for review.

**BREED**

Miniature Poodle

**RADIOGRAPHIC FINDINGS**

**Right Stifle**

**SEX**

Patient has a history of TPLO of the right stifle 2 years ago.

F

The implants are in situ and intact. No evidence of loosening or failure is seen.

**AGE**

A nodular soft tissue opacity is superimposed onto the soft tissues distal and lateral of the TPLO site.

8 Years

Mild articular swelling of the right stifle joint is seen. There is a moderate amount of periarticular osteophytes. The patella is in situ.

**INTERPRETED BY**

Nele Eley, DVM  
Dr. med. Vet. DipECVDI

**Left Stifle**

The patella is medially luxated in both orthogonal views. Mild periarticular osteophytes are seen as well as mild articular swelling.

**HOSPITAL NAME**

Paws and Claws Pet  
Care

**Pelvis**

The muscle volume of both hind limbs appears to be mildly reduced.

Both coxofemoral joints present moderate osteoarthritic changes as a consequence of a dysplastic conformation of the coxofemoral joints.

**REFERRING VET**

Dr. Juarez

**RADIOGRAPHIC DIAGNOSIS**

- History of TPLO of the right stifle with no radiographic evidence of complication other than moderate osteoarthritis.
- In situ position of the right patella during the radiographic examination.
- Medial luxation of the left patella during the radiographic examination.
- Mild osteoarthritis of the left stifle.
- Moderate bilateral coxofemoral osteoarthritis secondary to hip dysplasia.

**INVOICE**

47991

**DATE**

10-25-21



**PATIENT**

Ruby Morgan

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

No evidence of implant failure or loosening is seen. The findings of the right stifle are compatible with the history of cranial cruciate ligament injury and TPLO. The patella is in situ at least during the radiographic examination in the right stifle. Grade 3 or 4 medial patella luxation appears to be present in the left stifle. Clinical correlation is required.

**SPECIES**

Canine

Both coxofemoral joints present moderate osteoarthritic changes; left more than right, as a consequence of canine hip dysplasia. In small breed and lightweight dogs, this is not necessarily of clinical significance.

**BREED**

Miniature Poodle

**SEX**

F

**AGE**

8 Years

**INTERPRETED BY**

Nele Eley, DVM  
Dr. med. Vet. DipECVDI

**HOSPITAL NAME**

Paws and Claws Pet  
Care



**REFERRING VET**

Dr. Juarez

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

**INVOICE**

47991

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**DATE**

10-25-21

**Nele Eley, DVM, Dr. med. vet., DipECVDI**  
European Specialist in Veterinary Diagnostic Imaging, Cert. Radiology,  
Senior lecturer University of Giessen, Germany, Veterinary Faculty, Department of Radiology  
Nele.Eley@sonopath.com