



**PATIENT**

Cooper Al Maso

**PRESENTING CLINICAL SIGNS**

Limping and leg lifting on the right hind since few months, and occasionally on L hind. On physical exam there is a swelling in the hock joint in the same leg. Antibiotics and anti-inflammatory were given, the swelling went down but Cooper still limping. recently cooper has hemorrhages on the skin b/w attached below

**SPECIES**

Canine

**RADIOGRAPHIC STUDY OF THE STIFLES, PELVIS, & TARSI**

Multiple lateral and orthogonal views of the stifles and tarsi and ventrodorsal hip extended views of the pelvis totaling 13 images available for review.

**BREED**

Cockapoo

**RADIOGRAPHIC FINDINGS**

**Stifles**

Mild bilateral articular swelling of the stifle joints is noted without evidence of osteoarthritis or cranial thrust of the tibiae. The patellae are in situ in all available images.

**SEX**

Male Neutered

**Pelvis**

No significant muscle atrophy is seen in the hind limbs.

**AGE**

6 Years

The pelvic bones and coxofemoral joints present within age and breed related normal limits. There is no evidence of dysplasia or osteoarthritis. No traumatic osseous injury is seen.

**INTERPRETED BY**

Nele Eley, DVM  
Dr. med. Vet. DipECVDI

**Tarsi**

Mild right tibiotarsal joint swelling is noted as well as mild periarticular swelling of the soft tissues dorsal to the distal right tibia and tibiotarsal joint with no evidence of concurrent osseous injury. The tibiotarsal joint is in situ. The subchondral bone surfaces are intact.

**HOSPITAL NAME**

Truscott Animal  
Hospital

**RADIOGRAPHIC DIAGNOSIS**

- Mild bilateral articular swelling of the stifle joints.
- Mild articular and moderate periarticular swelling of the right tibiotarsal joint without evidence of concurrent osseous injury.

**REFERRING VET**

Medhat Meawad

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

The radiographic study reveals mild bilateral articular swelling of the stifle joints without radiographic evidence of osteoarthritis, osteochondrosis, or cranial cruciate ligament injury. The patellae are in situ.

**INVOICE**

52119

The right tibiotarsal joint presents mild articular swelling and periarticular swelling is seen dorsal and lateral of the right tibiotarsal joint.

**DATE**

5-11-22

Consider polyarthropathy such as immune mediated as well as soft tissue injury in the right hind limb such as inflammation, infection, or traumatic injury. Correlate with the clinical findings. Aspiration and analysis of synovia could be considered for further definition as well as ultrasound with eventual fine needle aspiration of the soft tissue swelling in the right tibiotarsal area.



**PATIENT**

Cooper Al Maso

**SPECIES**

Canine

**BREED**

Cockapoo

**SEX**

Male Neutered

**AGE**

6 Years

**INTERPRETED BY**

Nele Eley, DVM  
Dr. med. Vet. DipECVDI

**HOSPITAL NAME**

Truscott Animal  
Hospital

**REFERRING VET**


Medhat Meawad

**INVOICE**

52119

**DATE**

5-11-22



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Nele Eley**, DVM, Dr. med. vet., DipECVDI  
European Specialist in Veterinary Diagnostic Imaging, Cert. Radiology,  
Senior lecturer University of Giessen, Germany, Veterinary Faculty, Department of Radiology  
Nele.Eley@sonopath.com