

**PATIENT**

Ray Ciolkosz

PRESENTING CLINICAL SIGNS

acute onset swelling of left front leg, anorexia, fever

SPECIES

Canine

RADIOGRAPHIC STUDY OF THE LEFT FRONT LIMB & ABDOMEN

Mediolateral views of the left front limb and right/left lateral and ventrodorsal views of the abdomen totaling 6 images available for review.

BREED

Lab Mix

RADIOGRAPHIC FINDINGS**Abdomen**

The surrounding bony structures are within normal limits.

SEX

Male

No abnormalities of the extraabdominal soft tissues are noted. The abdominal wall is smooth and thin.

The liver is appropriate in position, size and presents uniform opacity.

AGE

4

The splenic head is in the anticipated position and within normal limits for size and opacity. The splenic body and tail are considered normal for position, size, shape and opacity.

Both kidneys are seen and present with normal size, shape, delineation and opacity. The urinary bladder is in its anticipated position. No radiopaque calculi are noted throughout the upper and lower urinary tract.

INTERPRETED BYNele Eley, DVM
Dr. med. Vet. DipECVDI

The stomach is post-prandial.

No evidence of abnormal dilation of the small intestinal loops is seen; however, one small intestinal segment appears to present narrow winding and luminal gas pattern with the impression of regionally reduced serosal detail. This is seen on the left lateral view only (see image below) and not repeated in the right lateral or orthogonal views.

HOSPITAL NAMERockaway Animal
Hospital

The large intestine contains a moderate amount of fecal material.

REFERRING VET

Dr. Maniar

Left Front Limb

The osseous structures of the left front limb present within normal limits.

The shoulder, elbow, and carpus present no deviation from their expected anatomy. There is no evidence of abnormal soft tissue swelling.

INVOICE

47563

The fascial planes of the forearm and around the humerus are well seen and delineated by thin, fat, opaque outline as expected.

DATE

9-27-21



PATIENT

Ray Ciolkosz

RADIOGRAPHIC DIAGNOSIS

- Segmental small intestinal maldigestion pattern with potential plication seen on the left lateral view only - otherwise, radiographically normal abdomen.
- Radiographically left front limb.

SPECIES

Canine

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The impression of segmental small intestinal plication is likely to be incidental; however, a minimal amount of gravel appears to be present in the same small intestinal segment and subileus cannot be ruled out. Based on the radiographic findings, further definition by means of abdominal ultrasound would be reasonable in order to rule out subileus and regional peritonitis. No definitive osseous and soft tissue pathology was evident in the radiographic study of the left front limb. However, the sensitivity of radiographs for soft tissue changes is low and correlation with the clinical findings is recommended.

BREED

Lab Mix

SEX

Male

AGE

4

INTERPRETED BY

Nele Eley, DVM
Dr. med. Vet. DipECVDI

HOSPITAL NAME

Rockaway Animal
Hospital

REFERRING VET

Dr. Maniar

INVOICE

47563

DATE

9-27-21





PATIENT

Ray Ciolkosz

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

SPECIES

Canine

Nele Eley, DVM, Dr. med. vet., DipECVDI
European Specialist in Veterinary Diagnostic Imaging, Cert. Radiology,
Senior lecturer University of Giessen, Germany, Veterinary Faculty, Department of Radiology
Nele.Eley@sonopath.com

BREED

Lab Mix

SEX

Male

AGE

4

INTERPRETED BY

Nele Eley, DVM
Dr. med. Vet. DipECVDI

HOSPITAL NAME

Rockaway Animal
Hospital

REFERRING VET

Dr. Maniar

INVOICE

47563

DATE

9-27-21