



PATIENT

Max Farlow

PRESENTING CLINICAL SIGNS

Upper airway congestion and nasal discharge.
Abnormal PE/Chem/CBC/UA Results: No nasal discharge noted prior nor during the scan

SPECIES

Feline

COMPUTED TOMOGRAPHIC STUDY OF THE HEAD

Plain and post contrast studies available for review.

BREED

DSH

COMPUTED TOMOGRAPHIC FINDINGS

The pictured parts of the dentition are complete and unremarkable in all jaw quadrants.

SEX

Male Neutered

The nasopharynx is obstructed by an ovoid well delineated soft tissue attenuating mass of approximately 13mm length, 13mm width, and 10mm height. Mild nasopharyngeal fluid accumulation is seen. There is no evidence of concurrent otitis. Minimal mucosal swelling is seen in the caudal aspect of the right nasal cavity.

AGE

12 Years, 5 Months

Both temporomandibular joints present congruent joint spaces with even subchondral bone surfaces and are considered within normal limits.

The tympanic bullae are aerated, the mucosal lining is not seen, and the bony wall is smooth and thin. The external ear canals are within normal limits.

INTERPRETED BY

Nele Eley, DVM
Dr. med. Vet. DipECVDI

The brain presents no deviation from normal anatomy and symmetry. The brain parenchyma is homogeneous and within normal limits for attenuation and distribution of contrast enhancement. The ventricular system is non-dilated and symmetric.

The submandibular and medial retropharyngeal lymph nodes are small and elongated with a normal short-to-long-axis-ratio of < 0.5. The attenuation and contrast enhancement pattern are uniform.

HOSPITAL NAME

Mobile Pet Imaging
CFL

COMPUTED TOMOGRAPHIC DIAGNOSIS

- Nasopharyngeal polyp - no evidence of concurrent otitis.

REFERRING VET

Rodriguez

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS


The CT study reveals upper airway obstruction by a nasopharyngeal polyp. The origin of the polyp remains unclear. It is the typical position for middle ear polyps secondary to chronic middle inflammation; however, at least at this point, no evidence of otitis media, externa, or interna is seen. Polyp removal is recommended regardless.

INVOICE

47596

DATE

9-29-21



PATIENT

Max Farlow

SPECIES

Feline

BREED

DSH

SEX

Male Neutered

AGE

12 Years, 5 Months

INTERPRETED BY

Nele Eley, DVM
Dr. med. Vet. DipECVDI

HOSPITAL NAME

Mobile Pet Imaging
CFL

REFERRING VET


Rodriguez


INVOICE

47596

DATE

9-29-21





PATIENT

Max Farlow

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

SPECIES

Feline

Nele Eley, DVM, Dr. med. vet., DipECVDI
European Specialist in Veterinary Diagnostic Imaging, Cert. Radiology,
Senior lecturer University of Giessen, Germany, Veterinary Faculty, Department of Radiology
Nele.Eley@sonopath.com

BREED

DSH

SEX

Male Neutered

AGE

12 Years, 5 Months

INTERPRETED BY

Nele Eley, DVM
Dr. med. Vet. DipECVDI

HOSPITAL NAME

Mobile Pet Imaging
CFL

REFERRING VET

Rodriguez

INVOICE

47596

DATE

9-29-21