



**PATIENT PRESENTING CLINICAL SIGNS**

**Cloey Balisky** History: urinary dribbling when asleep with normal micturition when awake, no other lower urinary signs described.

**SPECIES** Physical Examination: N/A.

Canine Urinalysis: SG 1.020, negative sediment.

**BREED** CBC: Pending.

Cockerpoo Serum Biochemistry: Pending.

Radiographic Findings: N/A.

**SEX**

FS

**AGE**

13 years

**WEIGHT**

8 kg

**INTERPRETED BY**

Remo Lobetti, BVSc,  
MMedVet (Med), PhD, Dipl.  
ECVIM

**ULTRASONOGRAPHIC EXAMINATION OF THE ABDOMEN**

**Urinary System**

Full urinary bladder with a normal appearance and thickness of the wall. Normal anechoic urine with no sediment or uroliths evident.

Normal trigone area, proximal urethra, and iliac blood vessels.

Normal iliac lymph nodes. Ureters not visualized.

Normal renal size with increased echogenic appearance, some loss of cortico-medullary differentiation, irregular capsule, normal blood flow, and bilateral pyelectasia (left worse than right).

**Reproductive System**

N/A.

**IMAGING PERFORMED BY**

Dr Matthew Olcha

**Adrenal Glands**

Normal shape, echogenic appearance, size, and position.

**HOSPITAL NAME**

East Meadow Veterinary  
Center

**Spleen**

Normal size and echogenic appearance. Smooth homogenous parenchyma, smooth curvi-linear capsule, and normal vasculature. No evidence of inflammatory, neoplastic, infarction, or infiltrative changes noted.

**REFERRING VET**

Dr Matthew Center

**Liver**

Normal size, echogenic, and portal markings. No nodules or masses evident. Small gall bladder containing normal anechoic bile. Normal thickness and echogenic appearance of the gall bladder wall. Normal bile duct.

**INVOICE**

303066

**Gastrointestinal**

**DATE**

6/25/22

Normal appearance of the stomach, duodenum, small intestine, ileo-cecal junction, and colon with no loss of layering, normal wall thickness and peristaltic activity, and no distension of the lumen.



**PATIENT** *Pancreas*

Cloey Balisky Normal size and echogenic appearance. Normal echogenic appearance of the mesentery and fat surrounding the pancreas.

**SPECIES** *Free Abdomen*

Canine No mesenteric lymphadenomegaly.  
No ascites.

**BREED**

Cockerpoo

**ULTRASONOGRAPHIC FINDINGS**

Primary Findings:

- Renal disease.

Secondary Findings:

- None.

**SEX**

FS

**AGE**

13 years

**WEIGHT**

8 kg

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

Etiologies for the kidneys would be age-related denegation, early chronic kidney disease, pyelonephritis, and bacterial nephritis.

**INTERPRETED BY**

If the pending blood work rules out renal disease (normal urea, creatinine, and SDMA) then urinary sphincter mechanism incompetence and spinal disease would be important considerations.

Initial further assessment that could be considered would be neurological examination.

Specific therapy would be dependent on an etiological diagnosis.

Remo Lobetti, BVSc,  
MMedVet (Med), PhD, Dipl.  
ECVIM

**IMAGING PERFORMED BY**

Dr Matthew Olcha

**HOSPITAL NAME**

East Meadow Veterinary  
Center

**REFERRING VET**

Dr Matthew Center

**INVOICE**

303066

**DATE**

6/25/22



**PATIENT IMAGES**

Cloey Balisky **Left kidney**

**SPECIES**

Canine

**BREED**

Cockerpool

**SEX**

FS

**AGE**

13 years

**WEIGHT**

8 kg

**INTERPRETED BY**

Remo Lobetti, BVSc,  
MMedVet (Med), PhD, Dipl.  
ECVIM

**IMAGING PERFORMED BY**

Dr Matthew Olcha

**HOSPITAL NAME**

East Meadow Veterinary  
Center

**REFERRING VET**

Dr Matthew Center

**INVOICE**

303066

**DATE**

6/25/22



**The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.**

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance, please contact me.

**Remo Lobetti, BVSc, MMedVet (Med), PhD, Dipl. ECVIM (Internal Medicine)**  
[rlobetti@mweb.co.za](mailto:rlobetti@mweb.co.za)