



**PATIENT**

Zoey Cirelli

**SPECIES**

Feline

**BREED**

Maine Coon Mix

**SEX**

Neutered Male

**AGE**

9 Years 1 Month

**WEIGHT**

19.6 Pounds

**INTERPRETED BY**

Sara Brethel,  
DVM, DACVIM  
(Cardiology)

**IMAGING PERFORMED BY**

Dr. Hesham Akbawy

**HOSPITAL NAME**

Lincoln Avenue Cat  
and Dog Hospital

**REFERRING VET**

Dr. Hesham Akbawy

**INVOICE**

37276

**DATE**

6/1/26

**PRESENTING CLINICAL SIGNS**

History: Has been on enalapril for a year, his BNP was 1100 dropped to 700ish

**ULTRASONOGRAPHIC EXAMINATION OF THE HEART**

FELINE CARDIAC PARAMETERS	BODY WEIGHT (kg)	HR (BPM)	IVSd (cm)	LVIDd (cm)	LVWd (cm)	FS (%)	EF (%)
<b>NORMAL PARAMETER</b>	-----	150-240	0.3-0.6	1.0-2.1	0.25-0.6	35-67	80-100
<b>PATIENT</b>	8.9	NM	0.55	1.47	0.76	31.97	--
FELINE CARDIAC PARAMETERS	LA/AO (M-mode)	LA/AO HEART BASE (Sisson)	LAD LA MAX 4 Chamber		LVOT VEL. (m/s)	RVOT VEL. (m/s)	IVRT (m/)
<b>NORMAL PARAMETER</b>	<1.5	1.6	0.7-1.7		<1.6	<1.3	40-60
<b>PATIENT</b>	--	--	--		~1.0	~0.5	NM

Adapted from June Boon, Veterinary Echocardiography, 1998  
Sisson D et al. JVIM 1991; 5: 232, Jacobs et al. Am J Vet Res 1985; 46:1705

LVIDs: 1.0 \*\* No other measurements were obtained.

**Cardiac Presentation**

The left atrium is within normal limits. The mitral valve leaflets are normal and there is no mitral regurgitation. There is no evidence of systolic anterior motion of the mitral valve and no evidence of a left ventricular outflow tract obstruction. There is concentric hypertrophy of the left ventricle. The right atrium is normal. The tricuspid valve is normal without evidence of tricuspid regurgitation. The right ventricle appears to have preserved systolic function subjectively. The aortic and pulmonic valves are normal without evidence of insufficiency. Aortic and pulmonic outflow velocities are within normal limits. The aorta and PA are normal along with the associated PA branches. There is no evidence of pleural effusion, pericardial effusion, or intracardiac masses.

**ULTRASONOGRAPHIC FINDINGS**

- Left ventricular concentric hypertrophy
- Normal left atrial size

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

The patient has evidence of left ventricular concentric hypertrophy and is classified as a stage B1 due to the normal left atrial size. If not already performed, it is recommended to ensure that patients blood pressure is normal and the patient is euthyroid. If the patient is euthyroid and normotensive, then the patient has underlying hypertrophic cardiomyopathy. No cardiac medications are indicated



## PATIENT

Zoey Cirelli

## SPECIES

Feline

## BREED

Maine Coon Mix

## SEX

Neutered Male

## AGE

9 Years 1 Month

## WEIGHT

19.6 Pounds

## INTERPRETED BY

Sara Brethel,  
DVM, DACVIM  
(Cardiology)

## IMAGING PERFORMED BY

Dr. Hesham Akbawy

## HOSPITAL NAME

Lincoln Avenue Cat  
and Dog Hospital

## REFERRING VET

Dr. Hesham Akbawy

## INVOICE

37276

## DATE

6/1/26

at this time as the patient is at a low risk for complications associated with this condition. Since this can be a progressive condition, serial monitoring is recommended. It's recommended to recheck an echocardiogram in 6 months, sooner if the patient develops cardiovascular clinical signs.

If the patient does not have a history of systemic hypertension or proteinuria, ACE inhibitor therapy does not appear to be indicated. With the normal left atrial size, typically cardiac therapies are not recommended at this time. Serial monitoring should be performed though, as this can be a progressive condition.

Recommend obtaining a blood pressure on the patient to ensure it is <160mmHg. If the blood pressure is elevated recommend following ACVIM guidelines for systemic hypertension and treating if indicated.

Standard perioperative fluid rates should be well-tolerated. Medications like dexmedetomidine and other alpha 2 agonists are best avoided. Ketamine is also best avoided. Anticholinergics can be used in the case of a clinically significant bradyarrhythmia (i.e., bradycardia with concurrent hypotension). If the patient is on an ACEi, recommend not giving this therapy the day of anesthesia.



**The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.**

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sara Brethel DVM, DACVIM (Cardiology)

[info@SonoPath.com](mailto:info@SonoPath.com)