



PATIENT

Crichton Hernandez

SPECIES

Canine

BREED

Poodle Mix

SEX

Male

AGE

2Y

WEIGHT

78

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet.
DipECVDI

IMAGING PERFORMED BY

JSS/JRD

HOSPITAL NAME

Parrish Creek
Veterinary Clinic

REFERRING VET

Dr. Daugherty

INVOICE

73311

DATE

1-13-26

PRESENTING CLINICAL SIGNS

Abdominal exploratory surgery for cryptorchid testicle for attempted castration (Unsuccessful).
Abdominal Ultrasound performed today, unable to locate testicle.

COMPUTED TOMOGRAPHY OF THE ABDOMEN

A high resolution pre- and post-contrast CT study of the abdomen is provided for review.

COMPUTED TOMOGRAPHIC FINDINGS

The hind limbs are positioned lateral to the abdomen, causing moderate image noise due to photon starvation.

The serosal fat presents normal attenuation behavior. There is no evidence of peritoneal effusion or peritonitis.

Both kidneys present within normal limits for size, shape and organ architecture. After contrast administration, a bilaterally symmetric and uniform nephro- and pyelogram is noted.

The left testicle is located in the scrotum. The right testicle cannot be appreciated.

The adrenal glands are within normal limits for size, shape and organ architecture.

Both liver and spleen present with normal shape, even surface, uniformly attenuating parenchyma and homogeneous contrast enhancement, unremarkable.

The pancreas is evenly contoured; the pancreatic parenchyma is homogeneous and presents uniform contrast enhancement.

The position, delineation, wall and content of the gastrointestinal tract are considered within normal limits throughout.

The bony and surrounding soft tissue structures reveal no abnormalities.

COMPUTED TOMOGRAPHIC DIAGNOSIS

- Right testicle cannot be visualized

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The right testicle cannot be identified, and the presumptive diagnosis can be testicular regression syndrome. If castration of the left testicle is considered, measurement of the anti-Müllerian hormone can be performed to check if there is retained functional gonadal tissue – there should be an interval of at least 14 days between castration and the measurement. If positive, consider a follow-up CT with hind limbs extended caudally to prevent photon starvation.



PATIENT

Crichton Hernandez

SPECIES

Canine

BREED

Poodle Mix

SEX

Male

AGE

2Y

WEIGHT

78

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet.
DipECVDI

IMAGING PERFORMED BY

JSS/JRD

HOSPITAL NAME

Parrish Creek
Veterinary Clinic

REFERRING VET

Dr. Daugherty

INVOICE

73311

DATE

1-13-26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
info@sonopath.com