



PATIENT PRESENTING CLINICAL SIGNS

Kiki Gorr Bone tumor left scapula, biopsy chondrogenic sarcoma ~3x3cm firm bony mass left scapula otherwise healthy exam Oncologist requested MET Check. Rx: +/- gabapentin

SPECIES COMPUTED TOMOGRAPHY OF THE SKULL, THORAX, AND ABDOMEN

Feline A high resolution pre- and post-contrast CT study of the thorax and abdomen and a post contrast CT study of the skull are provided for review.

BREED COMPUTED TOMOGRAPHIC FINDINGS

DLH Skull

A supernumerary triadan 107&207 is seen. Triadan 106, 307, 407 present with advanced tooth root resorption.

SEX

FS The nasal cavity presents the expected aerated spaces between thin & even conchae and turbinates with smooth mucosal lining.

AGE

Both temporomandibular joints present congruent joint spaces with even subchondral bone surfaces and are considered within normal limits.

6 Years

Both tympanic bullae are aerated, the mucosal lining is not seen, the bony wall is smooth and thin. The external ear canals are within normal limits.

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

The brain presents no deviation from normal anatomy and symmetry. The brain parenchyma is homogeneous and within normal limits for attenuation and distribution of contrast enhancement. The ventricular system is non-dilated and symmetric.

HOSPITAL NAME

Mobile Pet Imaging
CFL

The submandibular and medial retropharyngeal lymph nodes are small and elongated with a normal short-to-long-axis-ratio is < 0.5, the attenuation and contrast enhancement pattern is uniform.

Thorax

REFERRING VET

Borecky

Protruding from the distal segment of the left spina scapulae a well-defined, cauliflower like, heterogeneous mineralizing expansile mass with irregular margins is visible measuring 3.1 x 2.2 x 3.0 cm in size. The overlying cutaneous tissue presents a focal irregularity – likely history of biopsy.

INVOICE

49622

The sternal, cranial mediastinal and tracheobronchial lymph nodes are small elongated with a normal short-to-long-axis-ratio is < 0.5, the attenuation and contrast enhancement pattern is uniform and considered within normal limits.

The cardiovascular structures including the pulmonary vasculature are within normal limits.

DATE

1-17-22

The bronchial tree presents with regular branching and tapers uniformly towards the periphery as expected, the bronchial walls are thin and smooth. The bronchus-to-artery ratio is within normal limits.



PATIENT The lung parenchyma presents the expected architecture with regions of compression atelectasis of the ventral dependent aspects of the lung.

Kiki Gorr

Small incidental gas pockets are seen within the esophageal lumen, there is no evidence of abnormal dilation.

SPECIES

Abdomen

Feline

The serosal fat presents normal attenuation behavior. There is no evidence of peritoneal effusion or peritonitis.

BREED

DLH

Both kidneys present within normal limits for size, shape and organ architecture. After contrast administration a bilaterally symmetric and uniform nephro- and pyelogram is noted.

SEX

FS

The adrenal glands are within normal limits for size, shape and organ architecture. Both liver and spleen present with normal shape, even surface, uniformly attenuating parenchyma and homogeneous contrast enhancement, unremarkable.

The pancreas is evenly contoured, the pancreatic parenchyma is homogeneous and presents uniform contrast enhancement.

AGE

6 Years

The position, delineation, wall and content of the gastrointestinal tract are considered within normal limits throughout.

INTERPRETED BY

The bony and surrounding soft tissue structures reveal no abnormalities.

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

COMPUTED TOMOGRAPHIC DIAGNOSIS

- Monostotic semiaggressive osteoproliferative mass left spina scapulae
- Supernumerary tooth elements 107&207
- Dental resorptive lesions 106, 307&407
- No evidence of pulmonary metastatic disease
- Normal abdomen

HOSPITAL NAME

Mobile Pet Imaging
CFL

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

REFERRING VET

Borecky

The scapular mass is consistent with the history of primary osseous neoplasia, such as chondrosarcoma, osteochondrosarcoma. The chances of scapulectomy as limp sparing technique might be discussed with surgeon/oncologist.

INVOICE

49622

DATE

1-17-22



PATIENT

Kiki Gorr

SPECIES

Feline

BREED

DLH

SEX

FS

AGE

6 Years

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI



HOSPITAL NAME

Mobile Pet Imaging
CFL

REFERRING VET

Borecky

INVOICE

49622

DATE

1-17-22

The information and recommendations provided are based on the images presented by the referring veterinarian. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com