



## PATIENT

Bunny Mourey

## SPECIES

Canine

## BREED

Greyhound

## SEX

FS

## AGE

8

## WEIGHT

34.9kg

## INTERPRETED BY

Sebastian Schaub, DVM  
Dr. med. vet.  
DipECVDI

## IMAGING PERFORMED BY

MP

## HOSPITAL NAME

Green Dog Dental and  
Wellness

## REFERRING VET

Dr Alzate

## INVOICE

73042

## DATE

12-17-25

## PRESENTING CLINICAL SIGNS

1-2cm gingival mass on the buccal aspect of 408. Appears partially fluid filled and blood profusely with incisional biopsy

## COMPUTED TOMOGRAPHY OF THE SKULL

A high resolution plain CT study of the skull is provided for review.

## COMPUTED TOMOGRAPHIC FINDINGS

The tooth elements 101, 105, 106, 110, 201-203, 205, 206, 208, 210, 301-303, 307, 311, 407, 409-411 are absent. At the distal aspect of triadan 408, level with the alveolar crest of absent triadan 409, a uniform soft tissue attenuating gingival mass is seen, extending into the body of the mandible at the same level – presenting an ill-defined zone with moth eaten osteolysis of the mandible at the same level.

The nasal cavity presents the expected aerated spaces between thin & even conchae and turbinates with smooth mucosal lining.

Both temporomandibular joints present congruent joint spaces with even subchondral bone surfaces and are considered within normal limits.

Both tympanic bullae are aerated, the mucosal lining is not seen, the bony wall is smooth and thin. The external ear canals are within normal limits.

The right mandibular lymph nodes are prominent.

## COMPUTED TOMOGRAPHIC DIAGNOSIS

- Soft tissue mass level with absent triadan 409 with monostotic aggressive osteolysis of the body of the mandible at the same level
- Lymphadenopathy right mandibular lymph nodes
- Multiple absent teeth, see above

## INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The clinically appreciated mass at the right mandible – level with absent triadan 409 is highly suggestive for primary soft tissue neoplasia with osseous involvement. Differentials can include fibrosarcoma, melanoma, squamous cell carcinoma, odontogenic neoplasia, other. Biopsy has already been performed for specification and results are pending. Complete surgical resection via hemimandibulectomy is considered feasible; possible osteotomy lines are distal to triadan 408 and level with absent triadan 410/411.

Recommend FNA sampling of the right mandibular lymph nodes to differentiate between reactive lymphoid hyperplasia or metastatic disease.



## PATIENT

Bunny Mourey

## SPECIES

Canine

## BREED

Greyhound

## SEX

FS

## AGE

8

## WEIGHT

34.9kg

## INTERPRETED BY

Sebastian Schaub, DVM  
Dr. med. vet.  
DipECVDI

## IMAGING PERFORMED BY

MP

## HOSPITAL NAME

Green Dog Dental and  
Wellness

## REFERRING VET


Dr Alzate

## INVOICE

73042

## DATE

12-17-25



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Sebastian Schaub**, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI  
[info@sonopath.com](mailto:info@sonopath.com)