



PATIENT

Mia Amaya

PRESENTING CLINICAL SIGNS

Presented for evaluation of a right forelimb lameness. The right forelimb lameness was noticed several years ago during puppyhood. Lameness has been intermittent and appears after exercise; bouts of lameness typically lasts a few days. Within the last 1.5 months, hindlimb lameness was also noted after laying down - owners unsure which hindleg. Normal EDUD; No VDSC; Energetic. Mia current medications: Fish oil, CBD Activity:3 walks per day, 10-20 minutes each Walks are reduced due to lameness Mia is currently eating a raw diet, with no history of allergies. Mia is UTD on vaccines.

SPECIES

Canine

BREED

Bernese Mountain Dog

COMPUTED TOMOGRAPHY OF THE FRONT LIMBS

A pre- and post-contrast CT study of front limbs in a soft tissue and bone reconstruction are provided for review.

COMPUTED TOMOGRAPHIC FINDINGS

SEX

FS

The osseous and surrounding soft tissue structures of the shoulder joints are within normal limits. No abnormalities of the region of the brachial plexus are appreciated.

AGE

3 Years

The elbow joints present mild osteophyte new bone formation, R>L. The tip of the medial coronoid process of the right elbow joint is irregular marginated, three isolated mineralized bodies are not at the craniomedial aspect of the tip of the right medial coronoid process, with the largest measuring up to 4.5 x 2.8 x 3.0 mm in size; the smallest is measuring <0.5 mm.

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

The tip of the medial coronoid process of the left elbow joint is demarcated by a fissure line running from the craniomedial to laterocaudal direction into the radio-ulnar joint space. The demarcated tip of the medial coronoid process of the left elbow joint is measuring 3.1 x 4.3 x 2.8 mm.

HOSPITAL NAME

Animal Health Partners

COMPUTED TOMOGRAPHIC DIAGNOSIS

- Fragmented medial coronoid process (FCP) elbow joints bilaterally
- Mild degenerative osteoarthritis elbow joints, R>L
- Structural normal elbow joints

REFERRING VET

Dr. Jeffery Biskup

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The findings are consistent with mild chronic degenerative joint disease of both elbow joints due to a fragmented medial coronoid process.


INVOICE

50381

Arthroscopy is considered beneficial to revise the elbow joints and remove the fragments of the medial coronoid process and prevent further damage.

DATE

2-17-22



PATIENT

Mia Amaya

SPECIES

Canine

BREED

Bernese Mountain
Dog

SEX

FS

AGE

3 Years

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Animal Health
Partners

REFERRING VET


Dr. Jeffery Biskup

INVOICE

50381

DATE

2-17-22





PATIENT

Mia Amaya

SPECIES

Canine

BREED

Bernese Mountain
Dog

SEX


FS

AGE

3 Years

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI



HOSPITAL NAME

Animal Health
Partners

REFERRING VET

Dr. Jeffery Biskup

INVOICE

50381

DATE

2-17-22

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com