



PATIENT

Oslo Maynard

SPECIES

Canine

BREED

Australian Shepherd

SEX

Male Neutered

AGE

3Y, 3M

WEIGHT

28.4kg

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet.
DipECVDI

IMAGING PERFORMED BY

Emily Johnson

HOSPITAL NAME

Bluegrass Veterinary
Specialists

REFERRING VET

Olivia Stone

INVOICE

73939

DATE

2-25-26

PRESENTING CLINICAL SIGNS

- 1 yr history of occasional limping on RH.
- O initially noticed after playing at the park
- not lame all the time
- carprofen helped when trial was done
- rest did not help

COMPUTED TOMOGRAPHY OF THE PELVIS, STIFLE JOINTS AND HIND PAWS

A pre- and post-contrast CT study of the pelvis and hind limbs in a bone and soft tissue reconstruction is provided for review.

COMPUTED TOMOGRAPHIC FINDINGS

The osseous and soft tissue structures of the caudal lumbar spine and lumbosacral junction reveal no abnormalities.

The osseous and surrounding soft tissue structures of the pelvis are within normal limits. The periarticular bones of the left coxofemoral joints present mild osteophyte new bone formation.

The volume of the musculature of the right hind limb is mildly decreased.

The periarticular bones of the right stifle joint present mild to moderate osteophyte new bone formation. The right stifle joint presents a mild intracapsular soft tissue swelling, distorting the infrapatellar fat-pad cranially. The left stifle joint has smooth margins of the periarticular bones and reveals no intracapsular soft tissue swelling.

The periarticular bones of both tibiotarsal joints present very mild osteophyte new bone formation. The surrounding soft tissue structures of the tarsus bilaterally is unremarkable.

The osseous and surrounding soft tissue structures of the hind paws reveal no abnormalities.

COMPUTED TOMOGRAPHIC DIAGNOSIS

- Osteoarthritis right stifle joint
- Mild effusion right stifle joint
- Mild disuse atrophy musculature right hind limb
- Mild osteoarthritis left coxofemoral joint
- Unremarkable lumbosacral junction

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The degenerative joint disease of the right stifle joint is highly concerning for underlying pathology of the cranial cruciate ligament ± meniscal pathology. Preceding trauma or arthritis are differentials. A positive drawer sign or tibial compression test under general anesthesia will support the diagnosis – partial rupture of the cranial cruciate ligament and potential thickening of the synovial capsule can result in only mild instability.



PATIENT

Oslo Maynard

SPECIES

Canine

BREED

Australian Shepherd

SEX

Male Neutered

AGE

3Y, 3M

WEIGHT

28.4kg

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet.
DipECVDI

IMAGING PERFORMED BY

Emily Johnson

HOSPITAL NAME

Bluegrass Veterinary
Specialists

REFERRING VET

Olivia Stone

INVOICE

73939

DATE

2-25-26





PATIENT

Oslo Maynard

SPECIES

Canine

BREED

Australian Shepherd

SEX

Male Neutered

AGE

3Y, 3M

WEIGHT

28.4kg

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet.
DipECVDI

IMAGING PERFORMED BY

Emily Johnson

HOSPITAL NAME

Bluegrass Veterinary
Specialists

REFERRING VET

Olivia Stone

INVOICE

73939

DATE

2-25-26



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
info@sonopath.com