



PATIENT PRESENTING CLINICAL SIGNS

Yuki Kauzlick Patient presented for acute onset this morning of no interest in food, lethargic, ataxic, head bobbing. No possible toxin ingestion. CT performed to diagnose liver vs. neurological disease
Abnormal PE/Chem/CBC/UA Results: ALT 128

SPECIES COMPUTED TOMOGRAPHY OF THE SKULL, THORAX AND ABDOMEN

Canine A high resolution pre- and post-contrast CT study of the skull and abdomen and a post-contrast CT study of the thorax are provided for review.

BREED COMPUTED TOMOGRAPHIC FINDINGS

Yorkshire Terrier Skull

Triadan 105, 311 and 411 are absent.

SEX The nasal cavity presents the expected aerated spaces between thin & even conchae and turbinates with smooth mucosal lining.

FS Both temporomandibular joints present congruent joint spaces with even subchondral bone surfaces and are considered within normal limits.

AGE Both tympanic bullae are aerated, the mucosal lining is not seen, the bony wall is smooth and thin. The external ear canals are within normal limits.

9 Months The brain presents no deviation from normal anatomy and symmetry. The brain parenchyma is homogeneous and within normal limits for attenuation and distribution of contrast enhancement. The ventricular system is non-dilated and symmetric.

INTERPRETED BY Sebastian Schaub, DVM
Dr. med. vet. DipECVDI The submandibular and medial retropharyngeal lymph nodes are small and elongated with a normal short-to-long-axis-ratio is < 0.5, the attenuation and contrast enhancement pattern is uniform.

HOSPITAL NAME Thorax

Animal Emergency Hospital Deland The bony and surrounding soft tissue structures are within normal limits.

REFERRING VET The sternal, cranial mediastinal and tracheobronchial lymph nodes are small elongated with a normal short-to-long-axis-ratio is < 0.5, the attenuation and contrast enhancement pattern is uniform and considered within normal limits.

Dr.Schwanebeck The cardiovascular structures including the pulmonary vasculature are within normal limits.

INVOICE The bronchial tree presents with regular branching and tapers uniformly towards the periphery as expected, the bronchial walls are thin and smooth. The bronchus-to-artery ratio is within normal limits.

56930 The lung parenchyma presents the expected architecture and attenuation behavior.

DATE Small incidental gas pockets are seen within the esophageal lumen, there is no evidence of abnormal dilation.

2-26-23 Abdomen

The serosal fat presents normal attenuation behavior. There is no evidence of peritoneal effusion

**PATIENT**

or peritonitis.

Yuki Kauzlick

Both kidneys present within normal limits for size, shape and organ architecture. After contrast administration a bilaterally symmetric and uniform nephro- and pyelogram is noted.

SPECIES

The adrenal glands are within normal limits for size, shape and organ architecture.

Canine

Both liver and spleen present with normal shape, even surface, uniformly attenuating parenchyma and homogeneous contrast enhancement, unremarkable.

BREED

The portal vein presents a normal order of its tributary veins and intrahepatic branching. No abnormal vessel is noted inside and outside of the liver parenchyma.

Yorkshire Terrier

The pancreas is evenly contoured, the pancreatic parenchyma is homogeneous and presents uniform contrast enhancement.

SEX

The position, delineation, wall and content of the gastrointestinal tract are considered within normal limits throughout.

FS

The bony and surrounding soft tissue structures reveal no abnormalities.

COMPUTED TOMOGRAPHIC DIAGNOSIS**AGE**

9 Months

- Multiple absent teeth
- Normal brain
- No evidence of portosystemic shunting, neither intra- nor extrahepatic
- Normal thorax

INTERPRETED BYSebastian Schaub, DVM
Dr. med. vet. DipECVDI**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

In the present study of the brain there is no evidence of macromorphological disease, explaining the acute presenting clinical signs.

HOSPITAL NAMEAnimal Emergency
Hospital Deland

If not yet done so the workup should be complemented by examination of CSF and complete bloodwork to screen for brain disease that is not necessarily associated with structural changes of the brain parenchyma and rule out other systemic illness. In case of the strong clinical suspicion of structural intraparenchymal changes an MRI may be considered. If there is evidence of hepatic dysfunction, hepatic biopsy is recommended.

REFERRING VET


Dr.Schwanebeck

INVOICE

56930

DATE

2-26-23



PATIENT

Yuki Kauzlick

SPECIES

Canine

BREED

Yorkshire Terrier

SEX

FS

AGE

9 Months

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Animal Emergency
Hospital Deland

REFERRING VET



Dr.Schwanebeck

INVOICE

56930

DATE

2-26-23





PATIENT

Yuki Kauzlick

SPECIES

Canine

BREED

Yorkshire Terrier

SEX

FS

AGE

9 Months

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Animal Emergency
Hospital Deland

REFERRING VET

Dr.Schwanebeck

INVOICE

56930

DATE

2-26-23

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com