



PATIENT

Myla Rondeau

PRESENTING CLINICAL SIGNS

Limping on LHL for 2 days, no swelling/ no bruising on paw pad noted during PE, reactive when LHL stifle manipulated- Tibial thrust/cranial drawer- absent
Abnormal PE/Chem/CBC/UA Results: BW- stress hyperglycemia, rest unremarkable

SPECIES

Feline

RADIOGRAPHIC STUDY OF THE STIFLE JOINTS

Radiographs of the stifle joints in two orthogonal imaging planes are provided for review.

BREED

Domestic Longhair

RADIOGRAPHIC FINDINGS

The growth plates are still faintly visible.

Both stifle joints present smooth osseous margins and without signs of intracapsular soft tissue swelling. The medial fabella of both stifle joints can be appreciated as pinpoint mineralized body.

SEX

Female Spayed

RADIOGRAPHIC DIAGNOSIS

- Normal stifle joints

AGE

18 Months

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The radiographic study of the stifle joints presents without abnormalities. The small medial fabella is a normal anatomical variant.

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

St. Catherine's Animal
Hospital

REFERRING VET

Dr. Jui Gokhale

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

INVOICE

56680

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

DATE

2-9-23

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com