



PATIENT

Bailey Silverberg

PRESENTING CLINICAL SIGNS

Owner reports 2 week history of off and on forelimb lameness. Owner thinks both are affected but the right is worse. Lameness most notable after walks/activity. 2 days ago, patient was so painful she was nearly non-ambulatory.

SPECIES

Canine

Abnormal PE/Chem/CBC/UA Results: Unable to identify lameness on gait exam today. Pain on flexion of the left shoulder/elbow. No pain identified anywhere else including the neck. Temp 102.5. No other concerns of physical exam. X-rays were taken without sedation. Bloodwork is pending.

BREED

Lab

RADIOGRAPHIC STUDY OF THE SHOULDER; ELBOW AND CARPAL JOINTS

A complete set of radiographs of the front limbs is provided for review.

SEX

FS

RADIOGRAPHIC FINDINGS

Both shoulder joints present smooth osseous margins, and no abnormalities of the surrounding soft tissue structures are appreciated.

AGE

5 Years, 1 Month

The periarticular bones present moderate osteophyte new bone formation. The tip of the contour of the medial coronoid process of both elbow joints is ill-defined/presents a decreased opacity. Moderate sclerosis is seen at the base of the medial coronoid process of both elbow joints. At the caudodistal aspect of the medial epicondyle of the left elbow joint, a well-defined fusiform, mineralized body is seen.

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

The osseous and surrounding soft tissue structures of both carpal joints present without abnormalities.

HOSPITAL NAME

Gentle Doctor Animal
Hospital

RADIOGRAPHIC DIAGNOSIS

- Coronoid disease elbow joints bilaterally
- Mild to moderate degenerative osteoarthritis elbow joints bilaterally
- Flexor enthesopathy left elbow joint with dystrophic mineralization
- Normal shoulder joints bilaterally
- Normal carpal joints bilaterally

REFERRING VET

Dr. Adrienne Kanne

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The degenerative joint disease of both elbow joints is likely a sequela to underlying pathology of the medial coronoid process (e.g. FCP). A CT study of the elbow joints can be used as advanced imaging modality and evaluation of the medial coronoid process and potential surgical planning.

INVOICE

51025

DATE

3-18-22



PATIENT

Bailey Silverberg

SPECIES

Canine

BREED

Lab

SEX

FS

AGE

5 Years, 1 Month

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Gentle Doctor Animal
Hospital

REFERRING VET

Dr. Adrienne Kanne

INVOICE

51025

DATE

3-18-22





PATIENT

Bailey Silverberg

SPECIES

Canine

BREED

Lab

SEX

FS

AGE

5 Years, 1 Month

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Gentle Doctor Animal
Hospital

REFERRING VET

Dr. Adrienne Kanne

INVOICE

51025

DATE

3-18-22

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com