



**PATIENT PRESENTING CLINICAL SIGNS**

Tito Flashner History: Intrapelvic mass, lameness affecting Left hind.

**COMPUTED TOMOGRAPHIC STUDY OF THE PELVIS**

**SPECIES**

Canine A high resolution pre- and post-contrast CT study of the pelvis is provided for review.

**COMPUTED TOMOGRAPHIC FINDINGS**

**BREED**

Bernadoodle A granular mineralizing mass is extending through the left obturator foramen, measuring 5.7 x 2.7 x 4.6 cm in size; a small stalk like extension of the mass is coursing craniodorsally up to the level of the ventral aspect of S1. The left obturator foramen is mildly widened and the surrounding osseous margins present solid mild periosteal new bone formation. The mass is occupying approximately 60% of the cross-sectional area of the pelvic canal, the rectum is deviated to the right and distorted. The mass is deviating the left sciatic nerve laterally and dorsally.

**SEX**

Neutered Male The descending colon contains a significant amount of feces.

**AGE**

5 Years 9 Months Multiple small (<2 mm) mineral attenuating calculi are seen in the urinary bladder. In the subcutaneous tissue at the left dorsolateral aspect of L6/L7, an ovoid shaped soft tissue nodule is seen.

**INTERPRETED BY**

Sebastian Schaub, DVM  
Dr. med. vet. DipECVDI

**COMPUTED TOMOGRAPHIC DIAGNOSIS**

- Mineralizing mass centered on the left obturator foramen of the pelvis, secondary stenosis of the pelvic canal
- Secondary constipation
- Cystolithiasis without signs of obstruction
- Non-specific subcutaneous nodule left dorsolateral aspect L6/L7

**HOSPITAL NAME**

Mobile Pet Imaging

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

**REFERRING VET**

Dr. Meaux

The appreciated mass is consistent with either primary soft tissue neoplasia with dystrophic mineralization (e.g., fibrosarcoma, hemangiosarcoma) or primary osseous neoplasia (e.g., periosteal osteosarcoma, chondrosarcoma). Biopsy may be used for further differentiation. Mechanical interaction with the left coxofemoral joint and the sciatic nerve will explain the left hind limb lameness. Possible palliative surgical or radio-oncological treatment options can be discussed with surgeon and oncologist, based on biopsy results.

**INVOICE**

21424

**DATE**

3/2/23



**PATIENT**

Tito Flashner

**SPECIES**

Canine

**BREED**

Bernadoodle

**SEX**

Neutered Male

**AGE**

5 Years 9 Months

**INTERPRETED BY**

Sebastian Schaub, DVM  
Dr. med. vet. DipECVDI

**HOSPITAL NAME**

Mobile Pet Imaging

**REFERRING VET**


Dr. Meaux

**INVOICE**

21424

**DATE**

3/2/23



**The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.**

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Sebastian Schaub**, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI  
sebast.schaub@gmail.com



**PATIENT**

Tito Flashner

**SPECIES**

Canine

**BREED**

Bernadoodle

**SEX**

Neutered Male

**AGE**

5 Years 9 Months

**INTERPRETED BY**

Sebastian Schaub, DVM  
Dr. med. vet. DipECVDI

**HOSPITAL NAME**

Mobile Pet Imaging

**REFERRING VET**

Dr. Meaux

**INVOICE**

21424

**DATE**

3/2/23