



**PATIENT PRESENTING CLINICAL SIGNS**

Orion Gebhardt Preoperative scan prior to mass removal surgery  
 Abnormal PE/Chem/CBC/UA Results: 3/23/23 K 5.9, NA/K Ratio 25 all else wnl

**SPECIES COMPUTED TOMOGRAPHY OF THE THORAX AND ABDOMEN**

Canine A high resolution pre- and post-contrast CT study of the abdomen and a plain CT study of the thorax are provided for review.

**COMPUTED TOMOGRAPHIC FINDINGS**

**BREED** Thorax

Catahoula In the subcutaneous tissue at the left craniolateral abdominal wall, level with the 11<sup>th</sup>/12<sup>th</sup> left rib, a well-defined, soft tissue attenuating nodule is appreciated, measuring 15 mm in diameter.

**SEX** The sternal, cranial mediastinal and tracheobronchial lymph nodes are small elongated with a normal short-to-long-axis-ratio is < 0.5, the attenuation is uniform.

Male Neutered The cardiovascular structures including the pulmonary vasculature are within normal limits.

**AGE** The bronchial tree presents with regular branching and tapers uniformly towards the periphery as expected, the bronchial walls are thin and smooth. The bronchus-to-artery ratio is within normal limits.

3.5 Years The lung parenchyma presents the expected architecture and attenuation behavior with randomly distributed interspersed punctuate mineralization and dystelectasis of the caudodorsal aspect of the left caudal lung lobe.

**INTERPRETED BY**

Sebastian Schaub, DVM Dr. med. vet. DipECVDI Small incidental gas pockets are seen within the esophageal lumen, there is no evidence of abnormal dilation.

Abdomen

**HOSPITAL NAME** The serosal fat presents normal attenuation behavior. There is no evidence of peritoneal effusion or peritonitis.

Catskill Veterinary Services, PLLC Both kidneys present within normal limits for size, shape and organ architecture. After contrast administration a bilaterally symmetric and uniform nephro- and pyelogram is noted.

**REFERRING VET** The adrenal glands are within normal limits for size, shape and organ architecture.

Dr. Joseph D'Abbraccio Both liver and spleen present with normal shape, even surface, uniformly attenuating parenchyma and homogeneous contrast enhancement, unremarkable.

The pancreas is evenly contoured, the pancreatic parenchyma is homogeneous and presents uniform contrast enhancement.

**INVOICE**

57795 The position, delineation, wall and content of the gastrointestinal tract are considered within normal limits throughout.

**DATE**

4-13-23

In the subcutaneous tissue at the right craniolateral abdominal wall, level with the 11<sup>th</sup>/12<sup>th</sup> left rib, multiple small gas inclusions are appreciated and the cutaneous surface is irregular – the nodular lesion is not appreciated anymore.



**PATIENT**      **COMPUTED TOMOGRAPHIC DIAGNOSIS**

Orion Gebhardt

- History of excision of a subcutaneous soft tissue nodule left craniolateral abdominal wall
- Pulmonary osteomas

**SPECIES**      **INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

Canine

The CT study is negative for pulmonary metastatic disease. The abdomen presents without abnormalities.

**BREED**

Catahoula

**SEX**

Male Neutered

**AGE**

3.5 Years

**INTERPRETED BY**

Sebastian Schaub, DVM  
Dr. med. vet. DipECVDI

**HOSPITAL NAME**

Catskill Veterinary  
Services, PLLC

**REFERRING VET**

Dr. Joseph  
D'Abbraccio

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

**INVOICE**

57795

**Sebastian Schaub**, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI  
sebast.schaub@gmail.com

**DATE**

4-13-23

