



PATIENT PRESENTING CLINICAL SIGNS

Buns Emery History: Hx of right maxilla mass at 501-504: Papillary squamous cell carcinoma Rechecking after surgical removal 11/9/21.

SPECIES COMPUTED TOMOGRAPHIC STUDY OF THE SKULL

Canine A high resolution pre- and post-contrast CT study of the skull is provided for review.

COMPUTED TOMOGRAPHIC FINDINGS

BREED

French Bulldog Triadan 105, 305 & 405 are absent. Medial and caudal deviation of the alveolar crest and dental element of triadan 104 into the nasal cavity is noted. Triadan 104 is retained within the alveolar crest and moderate widening of the alveolar bone in the apical region is noted; the crown is covered by gingiva. A supernumerary triadan 201 is noted.

SEX

Neutered Male The nasal cavity presents the expected aerated spaces between thin & even conchae and turbinates with smooth mucosal lining.

AGE

23 Pounds Both temporomandibular joints present congruent joint spaces with even subchondral bone surfaces and are considered within normal limits.
Both tympanic bullae are aerated, the mucosal lining is not seen, the bony wall is smooth and thin. The external ear canals are within normal limits.

INTERPRETED BY

Sebastian Schaub, DVM Dr. med. vet. DipECVDI The brain presents no deviation from normal anatomy and symmetry. The brain parenchyma is homogeneous and within normal limits for attenuation and distribution of contrast enhancement. The ventricular system is non-dilated and symmetric.

HOSPITAL NAME

AMC of Mt. Pleasant The submandibular and medial retropharyngeal lymph nodes are small and elongated with a normal short-to-long-axis-ratio is < 0.5, the attenuation and contrast enhancement pattern is uniform.

COMPUTED TOMOGRAPHIC DIAGNOSIS

REFERRING VET

- History of excision of squamous cell carcinoma
- Aberrant position of triadan 104 with potential development of follicular cyst level with the crown
- Absent triadan 102, 305 and 405
- Supernumerary triadan 201

Brook Fenamore, VMD

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

INVOICE

14725 The aberrant position of triadan 104 with major parts located in the nasal cavity might be a sequela to abnormal development due to preceding surgery. There is evidence of widening of the alveolar crest level with the crown, concerning for secondary development of follicular cyst – removal of the retained triadan 104 is considered beneficial.

DATE

4/14/22



PATIENT

Buns Emery

No signs for biologically aggressive lesions are appreciated.

SPECIES

Canine

BREED


French Bulldog

SEX

Neutered Male

AGE

23 Pounds



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

INTERPRETED BY

Sebastian Schaub,
DVM Dr. med. vet.
DipECVDI

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com

HOSPITAL NAME

AMC of Mt. Pleasant

REFERRING VET

Brook Fenamore,
VMD

INVOICE

14725

DATE

4/14/22