



PATIENT PRESENTING CLINICAL SIGNS

Dakota Katz This morning he didn't eat all his breakfast he only ate half of his food and that is not normal for him. As the day went on he started to get lethargic and his fecal was really runny. He was out in the yard playing this weekend. Was given a Bully stick on Saturday.

SPECIES Abnormal PE/Chem/CBC/UA Results: pe: pale sclera chocolate pudding diarrhea in room. Nails long. Quiet, quick to lay down bw: chem Chloride 107mmol/L

Canine

RADIOGRAPHIC STUDY OF THE ABDOMEN

Radiographs of the abdomen in three imaging planes are provided for review.

BREED

Labrador Retriever

RADIOGRAPHIC FINDINGS

The surrounding bony structures are within normal limits.

No abnormalities of the extraabdominal soft tissues are noted. The abdominal wall is smooth and thin.

SEX

The serosal detail is maintained throughout the peritoneal and retroperitoneal space.

Neutered Male

The liver is appropriate in position, size and presents uniform opacity.

AGE

The splenic head is in the anticipated position and within normal limits for size and opacity. The splenic body and tail are considered normal for position, size, shape and opacity.

2 Years, 1 Month

INTERPRETED BY

Both kidneys are seen and present with normal size, shape, delineation and opacity. The urinary bladder is in its anticipated position. No radiopaque calculi are noted throughout the upper and lower urinary tract.

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

The stomach is in its anticipated position and presents normal content. A small amount of mineral opaque material is seen in the pyloric region of the stomach.

HOSPITAL NAME

Elizabeth Animal
Hospital

The small intestinal loops are of even diameter and non-dilated, a small amount of chymus and gas is seen within the small intestinal loops.

The colon is empty and collapsed.

RADIOGRAPHIC DIAGNOSIS

- Empty colon

REFERRING VET

Leon Anderson, DVM

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

The empty colon is considered as a sequela to the history of diarrhea. The stomach still contains ingesta, regarding the history last food was given in the morning, suggestive for functional gastric emptying disorder. In combination with the history, the findings are most suggestive for underlying gastroenteritis. If clinical signs are refractory to therapy, recommend follow up radiographs or a complete abdominal ultrasound examination.

INVOICE

57835

DATE

4-17-23



PATIENT

Dakota Katz

SPECIES

Canine

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

BREED

Labrador Retriever

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com

SEX

Neutered Male

AGE

2 Years, 1 Month

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Elizabeth Animal
Hospital

REFERRING VET

Leon Anderson, DVM

INVOICE

57835

DATE

4-17-23