



PATIENT

Fergus Love

PRESENTING CLINICAL SIGNS

LFL lameness initially noticed January 2022 with no known trauma, rads WNL Feb 2022. Lameness worse after rest or heavy activity. Lameness is noticed daily. No progression of lameness. Did not improve with gabapentin, and Metacam. Fergus also seeing a chiropractor (last 3 weeks ago). Fergus still going on walks and activity. Normal EDUD; No VDSC

SPECIES

Canine

COMPUTED TOMOGRAPHY OF THE FRONT LIMBS

A pre- and post-contrast CT study of front limbs in a bone and soft tissue reconstruction is provided for review.

BREED

English Springer Spaniel

COMPUTED TOMOGRAPHIC FINDINGS

The osseous and surrounding soft tissue structures of both front limbs present without abnormalities.

SEX

MN

The cranial tip of the medial coronoid process of the left elbow joint is irregular and a fissure line is demarcating an osseous fragment, measuring 3.4 x 2.8 x 3.1 mm in size. At the distal aspect of the tip of the medial coronoid process of the left elbow joint a secondary mineral attenuating fragment is seen, measuring 1.8 x 1.3 x 1.2 mm in size. Very mild osteophyte new bone formation is seen at the medial aspect of the left medial humeral condyle.

AGE

10 Months

The right elbow joint presents smooth osseous margins, the medial coronoid process is well defined, unremarkable.

The osseous and soft tissue structures of the carpal joints are within normal limits.

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

Post contrast administration no pathological contrast enhancement pattern is appreciated.

COMPUTED TOMOGRAPHIC DIAGNOSIS

- Fragmented medial coronoid process (FCP) left elbow joint
- Very mild degenerative osteoarthritis left elbow joint
- Structural normal shoulder joints
- Structural normal right elbow joint
- Structural normal carpal joints

HOSPITAL NAME

Animal Health Partners

INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS

REFERRING VET

Dr. Lea Mehrkens

The findings are consistent with very mild degenerative joint disease of the left elbow joint due to a fragmented medial coronoid process.

Arthroscopy is recommended to revise the left elbow joint and remove the fragments of the medial coronoid process and prevent further damage. Overall prognosis for the patient to be free of lameness is considered good after fragment removal. However, episodes of lameness may reoccur after exercise.

INVOICE

52902

DATE

7-15-22



PATIENT

Fergus Love

SPECIES

Canine

BREED

English Springer
Spaniel

SEX

MN

AGE

10 Months

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Animal Health
Partners

REFERRING VET


Dr. Lea Mehrkens

INVOICE

52902

DATE

7-15-22





PATIENT

Fergus Love

SPECIES

Canine

BREED

English Springer
Spaniel

SEX

MN

AGE

10 Months

INTERPRETED BY

Sebastian Schaub, DVM
Dr. med. vet. DipECVDI

HOSPITAL NAME

Animal Health
Partners

REFERRING VET


Dr. Lea Mehrkens

INVOICE

52902

DATE

7-15-22



The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

Sebastian Schaub, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI
sebast.schaub@gmail.com