



**PATIENT**

Finn Toth

**PRESENTING CLINICAL SIGNS**

Pet has had chronic lameness in the right forelimb. Was sent to us for second opinion of suspected right shoulder OCD, but on physical exam there was resistance to flexion of both elbows, degenerative changes in both elbows on radiograph, and no obvious OCD on lateral shoulder radiographs.

**SPECIES**

Canine

**COMPUTED TOMOGRAPHY OF THE FRONT LIMBS**

A high resolution plain and positive contrast athrography CT study of the front limbs is provided for review.

**BREED**

Golden Retriever

**COMPUTED TOMOGRAPHIC FINDINGS**

The osseous and surrounding soft tissue structures of the shoulder are within normal limits.

**SEX**

Male

The periarticular bones of both elbow joints present moderate osteophyte new bone formation. The tip of the medial coronoid process of both elbow joints is irregular margined and a fissure line is demarcating and osseous fragment respectively. The isolated osseous fragment is measuring 4.3 x 2.8 x 2.7 mm in size (R) and 4.1 x 3.9 x 2.9 mm (L). Both elbow joints present an incongruent joint space due to radio-ulnar step formation measuring 2.3 mm. Post contrast administration, the synovial lining of both elbow joints is prominent

**AGE**

10 Months

**COMPUTED TOMOGRAPHIC DIAGNOSIS**

- Fragmented medial coronoid process (FCP) elbow joints bilaterally
- Mild to moderate radio-ulnar step formation
- Moderate degenerative osteoarthritis elbow joints bilaterally
- Synovialitis elbow joints bilaterally
- Structural normal shoulder joints

**INTERPRETED BY**

Sebastian Schaub, DVM  
Dr. med. vet. DipECVDI

**HOSPITAL NAME**

Wilson Veterinary Hospital

**INTERPRETATION OF THE FINDINGS & FURTHER RECOMMENDATIONS**

The findings are consistent with moderate chronic degenerative joint disease of both elbow joints due to a fragmented medial coronoid process.

Arthroscopy is recommended to revise the elbow joints and remove the fragments of the medial coronoid process and prevent further damage.

**REFERRING VET**

Dr. Burge

Due to the radio-ulnar step formation, the chances of combining arthroscopy with dynamic double oblique proximal ulna-osteotomy should be discussed with orthopedic surgeon.

**INVOICE**

52944

**DATE**

7-18-22



**PATIENT**

Finn Toth

**SPECIES**

Canine

**BREED**

Golden Retriever

**SEX**

Male

**AGE**

10 Months

**INTERPRETED BY**

Sebastian Schaub, DVM  
Dr. med. vet. DipECVDI

**HOSPITAL NAME**

Wilson Veterinary  
Hospital

**REFERRING VET**


Dr. Burge

**INVOICE**

52944

**DATE**

7-18-22





**PATIENT**

Finn Toth

**SPECIES**

Canine

**BREED**

Golden Retriever

**SEX**


Male

**AGE**

10 Months

**INTERPRETED BY**

Sebastian Schaub, DVM  
Dr. med. vet. DipECVDI



**HOSPITAL NAME**

Wilson Veterinary  
Hospital

**REFERRING VET**

Dr. Burge

**INVOICE**

52944

**DATE**

7-18-22

The information and recommendations provided are based on the images presented by the referring veterinarian/sonographer. No evaluation can be communicated regarding pathology that was not visible in the image/video clips provided.

Thank you for this referral. If the clinical or image interpretation does not parallel your findings or if I can be of any further assistance please contact me.

**Sebastian Schaub**, Sebastian Schaub, DVM, Dr. med. vet. DipECVDI  
sebast.schaub@gmail.com